

#### N04/4/CHEMI/SP3/ENG/TZ0/XX



CHEMISTRY STANDARD LEVEL PAPER 3	School code
Thursday 18 November 2004 (morning)	Candidate code
1 hour	

#### INSTRUCTIONS TO CANDIDATES

1 hour

- Write your school code and candidate code in the boxes above.
- Do not open this examination paper until instructed to do so.
- Answer all of the questions from two of the Options in the spaces provided. You may continue your answers on answer sheets. Write your school code and candidate code on each answer sheet, and attach them to this examination paper using the tag provided.
- At the end of the examination, indicate the letters of the Options answered in the candidate box on your cover sheet and indicate the number of answer sheets used in the appropriate box on your cover sheet.

8804-6106 23 pages

(This question continues on the following page)

## Option A – Higher physical organic chemistry

A1.		There are two alcohols, $\bf A$ and $\bf B$ , with molecular formula $C_3H_8O$ . The following information was obtained from a mass spectrum of each alcohol.					
	<b>A</b> :	peaks at $m/z = 29, 31, 60$					
	<b>B</b> :	peaks at $m/z = 45, 60$					
	(a)	Write the formula for the species responsible for the peak at $m/z = 60$ .	[1]				
	(b)	Deduce the formula of the species with $m/z = 31$ .	[1]				
	(c)	Deduce the structures of the <b>two</b> alcohols.  Structure of <b>A</b>	[2]				
		Structure of <b>B</b>					
	(d)	The peak at $m/z = 45$ is more prominent than that at $m/z = 29$ . Suggest a reason for this.	[1]				

8804-6106

### (Question A1 continued)

(e)		e <sup>1</sup> H NMR spectrum of one of the alcohols shows four peaks with areas in the ratio 2:2:1.		
	(i)	State what can be deduced from this information.	[2	
		Four peaks		
		Areas in ratio 3:2:2:1		
	(ii)	Predict the number of peaks, and the ratio of their areas, in the <sup>1</sup> H NMR spectrum of the other alcohol.	[2	

**A2.** Gaseous hydrogen iodide, HI, decomposes into its elements when heated. The decomposition was investigated in a series of experiments carried out at the same temperature. The following data was obtained.

Experiment number	Initial [HI] / mol dm <sup>-3</sup>	Initial rate of reaction / mol dm <sup>-3</sup> s <sup>-1</sup>
1	$2.2 \times 10^{-3}$	1.1×10 <sup>-6</sup>
2	6.6×10 <sup>-3</sup>	9.9×10 <sup>-6</sup>
3	2.2×10 <sup>-2</sup>	1.1×10 <sup>-4</sup>
4	$4.4 \times 10^{-3}$	to be determined

(a)	Write the equation for the decomposition of hydrogen iodide.	[1]
(b)	Deduce the order of the reaction and explain your answer.	[2]
(c)	State the rate expression for the reaction.	[1]
(d)	Determine the initial rate of reaction in experiment 4.	[1]
(e)	Define the term <i>molecularity</i> and deduce its value in this reaction.	[2]

13.		formula and $pK_a$ value of chloroethanoic acid appear in Table 16 of the Data Booklet. Use	
	this i	nformation to answer the following questions.	
	(a)	Write the equation for the dissociation of chloroethanoic acid in aqueous solution.	[1]
	(b)	Deduce the $K_a$ expression for the dissociation.	[1]
	(c)	Calculate the value of $K_a$ for chloroethanoic acid.	[1]
	(d)	Arrange the following acids in increasing order of acid strength (starting with the weakest).	[1]
		chloroethanoic acid ethanoic acid iodoethanoic acid	
		Order	

## Option B – Medicines and drugs

B1.	Depressants such as tranquilizers and sedatives are capable of affecting the central nervous system.				
	(a)	State two effects, in each case, on the body of taking			
		(i)	a low dose of a tranquilizer.	[2]	
		(ii)	a high dose of a sedative.	[2]	
	(b)	Explain why depressants are sometimes described as anti-depressants.		[1]	
	(c)		most widely-used depressant is ethanol. Discuss the harmful effects of regularly taking amounts of ethanol by referring to <b>four</b> specific problems.	[4]	

## (Question B1 continued)

(d)	List two depressants whose structures are shown in Table 21 of the Data Booklet.				
(e)	One problem with many drugs is that users develop <i>tolerance</i> . Explain what is meant by the term <i>tolerance</i> and state why it could increase the risk to the user.	[2]			

B2.	Caffeine and nicotine are two stimulants whose structures are shown in Table 21 of the Data Booklet.				
	(a)	Describe <b>two</b> similarities in their structures, not including the presence of double bonds, methyl groups and nitrogen atoms.	[2]		
	(b)	Discuss the problems associated with nicotine consumption, distinguishing between short-term and long-term effects.	[6]		

Blank page

## Option C – Human biochemistry

C1.	The structures of the amino acids cysteine and serine are shown in Table 20 of the Data Boo They can react with each other to form a dipeptide.				
	(a)	State the type of reaction occurring when amino acids react together and identify the other product of the reaction.	[2]		
	(b)	Draw the structures of the <b>two</b> possible dipeptides formed in the reaction between one molecule of each of cysteine and serine.	[2]		
	(c)	Six tripeptides can be formed by reacting together one molecule of each of the amino acids arginine, histidine and leucine. Predict the primary structures of these six tripeptides using the symbols shown in Table 20 of the Data Booklet to represent the amino acids.	[3]		

(This question continues on the following page)

(d)		n many amino acid molecules react together a protein is formed. These proteins have ary, secondary and tertiary structures.		
	(i)	State the type of intermolecular force responsible for maintaining the secondary structure.	[1]	
	(ii)	State <b>two</b> other ways in which the tertiary structure of the protein is maintained.	[2]	

C2.	(a)	State	the empirical formula of all monosaccharides.	[1]
	(b)	The	structural formula of lactose is shown in Table 22 of the Data Booklet.	
		(i)	Deduce the structural formula of <b>one</b> of the monosaccharides that reacts to form lactose and state its name.	[2]
		(ii)	State the name of the <b>other</b> monosaccharide.	[1]
	(c)	State	e <b>two</b> major functions of polysaccharides in the body.	[2]

<b>3.</b>	3. The structural formulas of cholesterol and testosterone are shown in Table 22 of the Data Booklet.			
	(a)	Identify the class of compound to which cholesterol and testosterone belong.	[1]	
	(b)	State the names of <b>two</b> functional groups present in both cholesterol and testosterone.	[2]	
	(c)	Cholesterol and testosterone both contain a five-membered ring as part of their structures. Deduce the total number of hydrogen atoms joined directly to the carbon atoms in this ring.	[1]	

## **Option D – Environmental chemistry**

D1.	Part of fu	iculates are a type of primary air pollutant produced in several industries and by the burning uels.				
	(a)	The emission of particulates by some industries is reduced by an electrostatic method. Explain how this is done.	[3]			
	(b)	State <b>one</b> type of fuel that is very likely to produce particulates when burned.	[1]			
	(c)	Deduce the equation for a combustion reaction of methane in which particulates are formed.	[1]			

<b>D2.</b>	(a)	Exp	lain, with the help of an equation, why rain is naturally acidic.	[2]
	(b)	Cata	alytic converters are used in motor vehicles to reduce the emissions of acidic gases.	
		(i)	Give an equation to show the formation of nitrogen(II) oxide in a motor vehicle and identify the acid it forms in the atmosphere.	[2]
		(ii)	Nitrogen(II) oxide reacts with carbon monoxide in a catalytic converter to produce harmless substances. Deduce the equation for this reaction.	[2]

D3.	(a)	State what is meant by the term biological oxygen demand (BOD).	[2]
	(b)	Organic matter in water can be decomposed by both aerobic and anaerobic bacteria.	
		(i) State which type of bacteria is more likely to be active in water with a low BOD value.	[1]
		(ii) The following elements can occur in organic matter. Suggest <b>one</b> gas that is likely to be produced from each element when organic matter is decomposed by anaerobic bacteria.	[3]
		carbon	
		nitrogen	
		sulfur	
	(c)	Power stations may use river water for cooling purposes. Discuss the effects of this on fish in such a river.	[3]

## **Option E – Chemical industries**

£1.		extraction of metals from their ores often begins by using water to separate the ores from other erials found in the rock.	
	(a)	State what is done to the rock before water is used.	[1]
	(b)	An old method of extracting gold from its ore is <i>panning</i> , in which the gold and other impurities are mixed vigorously with water in a large dish. Suggest <b>one</b> physical property of gold that is important in its separation from impurities.	[1]
	(c)	A more modern method of extracting ores from rock uses <i>froth flotation</i> . Outline this method by referring to the substances used, apart from the ore and water, and stating how the ore is separated from the rock.	[3]
	(d)	Explain why the percentage of gold in the Earth's crust occurring as the element is much higher than that of iron.	[1]

E2.		Aluminium and iron are extracted from their ores by different chemical methods. For aluminium, electrolysis is used.					
	(a)	(i)	Identify the compound from which most aluminium is extracted.	[1]			
		(ii)	Electrolysis of this compound gives aluminium and another product. Write a half-equation for the formation of each product.	[3]			
	(b)	Mos	t iron is produced by heating iron ore with coke in a blast furnace.				
		(i)	State <b>two</b> other raw materials used in the blast furnace.	[1]			
		(ii)	Give the equation for the reduction of iron(III) oxide in the blast furnace.	[2]			

(a)		uss the differences between these <b>two</b> forms by referring to the amount of branching, the es between the polymer chains and the physical properties.
(b)	Both	forms of polythene are described as thermoplastics.
(b)	Both (i)	forms of polythene are described as <i>thermoplastics</i> .  State the meaning of this term.
(b)		
(b)		
(b)		State the meaning of this term.
(b)	(i)	State the meaning of this term.  Phenol-methanal is not a thermoplastic. State what type of polymer it is and how its

## Option F – Fuels and energy

a)	Outline how crude oil w	as formed.		
o)	Determine a value for the	$_{4}(g) + 9\frac{1}{2}O_{2}(g) \rightarrow 6CC$	$O_2(g) + 7H_2O(g)$	
))	$C_6H_1$	$_{4}(g) + 9\frac{1}{2}O_{2}(g) \rightarrow 6CC$	$O_2(g) + 7H_2O(g)$	
o)	C <sub>6</sub> H <sub>1</sub> Determine a value for the formation data.	$_{4}(g)+9\frac{1}{2}O_{2}(g)\rightarrow 6CC$ ne enthalpy of combus	$O_2(g) + 7H_2O(g)$ tion of hexane using th	ne following enthalpy
0)	$C_6H_1$ Determine a value for the formation data.  Compound	$_{4}(g) + 9\frac{1}{2}O_{2}(g) \rightarrow 6CC$ The enthalpy of combus $C_{6}H_{14}(g)$	$O_2(g) + 7H_2O(g)$ tion of hexane using the $CO_2(g)$	the following enthalpy $H_2O(g)$

(This question continues on the following page)

### (Question F1 continued)

(c)	A student burned some hexane in a calorimeter and used the data obtained to calculate a value for the enthalpy of combustion of hexane. Explain why this value was a lot smaller than the one obtained using the method in part (b).	[1]
(d)	Hexane is not suitable for use as a fuel in car engines because of its low octane number. Suggest why its isomer, 2,3-dimethylbutane, has a higher octane number.	[1]
		L-J

Turn over 8804-6106

(This question continues on the following page)

F2.	(a)	Outline <b>two</b> features of chemical reactions that do <b>not</b> apply to nuclear reactions.	[2]
	(b)	The isotope $^{218}Po$ can undergo either $\alpha\text{-decay}$ or $\beta\text{-decay}$ . Deduce the symbol and mass number of the element formed in each case.	[2]
		α-decay	
		β-decay	
	(c)	The half-life for the decay of <sup>218</sup> Po is 3.0 minutes. Calculate the mass of <sup>218</sup> Po remaining after a 12 g sample is left for 12 minutes.	[2]

8804-6106

# (Question F2 continued)

(d)	Discuss the concerns about safety in nuclear power plants by commenting on <b>two</b> aspects of <b>each</b> of the following.	
	<ul> <li>the withdrawal of all the control rods from the core</li> <li>the presence of oxygen in the coolant gas passing through the graphite moderator</li> <li>the breakage of a pipe carrying molten sodium in a breeder reactor.</li> </ul>	[6]