



CHEMISTRY STANDARD LEVEL PAPER 3

Tuesday 8 November 2011 (morning)

1 hour

| | C | andio | Jale | sessi | OH H | umb | er | |
|---|---|-------|------|-------|------|-----|----|--|
| 0 | 0 | | | | | | | |

Examination code

| 8 8 1 1 1 - 6 1 1 0 6 |
|-----------------------|
|-----------------------|

INSTRUCTIONS TO CANDIDATES

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- Answer all of the questions from two of the Options.
- Write your answers in the boxes provided.

Option A — Modern analytical chemistry

(a)

| A1. | Spectroscopy | involves | techniques | that | can | identify | substances | and | determine | their |
|------------|----------------|----------|------------|------|-----|----------|------------|-----|-----------|-------|
| | concentrations | 5. | | | | | | | | |

Identify the atomic or molecular processes associated with the UV and microwave

| | regions of the electromagnetic spectrum. | [2] |
|-----|--|-----|
| | UV: | |
| | | |
| | Microwave: | |
| | | |
| | | |
| (b) | Atomic absorption (AA) spectroscopy is used for the detection of metal ions. (i) Describe three principles of AA spectroscopy. | [3] |
| (b) | | [3] |

(This question continues on the following page)



[1]

(Question A1 continued)

(ii) Besides the detection of calcium ions in blood, water or soil samples, state **two** other uses of AA spectroscopy. Your answer should list the detection of two different metal ions, each from a different source.

| i) | lSi | _ | | | | | | | | | | | | ar | 1 (| d€ | €to | er | 'n | 11 | n | e | th | ie | C | 0 | n | ce | nt | ra | ıti | 0 | n | 0 | f a | 1 : | m | et | al | i | Эn | ιi | n | S | ol ¹ | ut | io | n | |
|----|-----|---|---|---|---|---|---|-------|---|---|-------|---|---|----|-----|----|-----|----|----|-----|---|---|----|----|---|---|---|----|----|----|-----|---|---|---|-----|-----|---|----|----|---|----|----|---|------------|-----------------|----|----|---|--|
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | • | • | • | • | ٠ | • | • | • | • | • | ٠ | • | | • | ٠ | ٠ | • | | · • | • | • | • | | • | • | | • | • | | • | • | | • | • | | ٠ | | • | | | ٠ | | . • | • | | | | |
| | | | | | • | | | | | • | • | | | | | | | | | | | | • | | | • | | • | • | | • | • | | • | | | | | | | | | | | • | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | . . | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | . - | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

| • | (a) | One type of molecular vibration that occurs when ${\rm CO_2}$ molecules are exposed to IR radiation is illustrated in the diagram below. | |
|---|-----|--|-----|
| | | → ← → | |
| | | O=C=O | |
| | | asymmetric stretching | |
| | | Identify \mathbf{two} other types of molecular vibrations that occur when CO_2 molecules are exposed to IR radiation. Illustrate your answer with appropriate diagrams. | [2] |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| (| (b) | Explain which molecular vibration of CO ₂ in (a) above is IR active and which is IR inactive . | [3] |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |



Nuclear magnetic resonance (NMR) and mass spectrometry are diagnostic techniques often

used in the identification of organic compounds. Deduce two similarities and one difference in the ¹HNMR spectra of the two (a) isomers CH₃COOH, a carboxylic acid, and HCOOCH₃, an ester. ¹H NMR data are given in Table 18 of the Data Booklet. [3] Similarities: Difference: The mass spectrum of one of the two isomers above has significant peaks at mass to charge ratios of 15, 45 and 60, while the other isomer has peaks at 15, 29, 31 and 60. Analyse these fragmentation patterns in the two mass spectra in order to distinguish between the two isomers. [2] Outline how the technique of magnetic resonance imaging (MRI) is used in body scanners. [2] (c)



Option B — Human biochemistry

B1. A student carried out an experiment to determine the energy value of 100.00 g of a food product by burning some of it. A 5.00 g sample was burned and the heat produced was used to heat water in a glass beaker.

She recorded the following data:

Mass of water heated = 100.00 g

Initial temperature of water = $19.2 \,^{\circ}$ C

Highest temperature of water = $28.6 \,^{\circ}$ C

Other data:

Heat capacity of the glass beaker = 90.2 J K^{-1}

Specific heat capacity of water = $4.18 \text{ J g}^{-1} \text{ K}^{-1}$

Calculate the energy value for 100.00 g of the food product, in kJ, showing your working. [3]

| | | |
|-------|------|--|
| | | |
| , | | |
| | | |
| | | |
| | | |



Proteins are macromolecules formed from 2-amino acids. Once a protein has been hydrolysed,

B2.

chromatography and electrophoresis can be used to identify the amino acids present. State the name of the linkage that is broken during the hydrolysis of a protein and (a) draw its structure. [2] Explain how electrophoresis is used to analyse a protein. [4] **B3.** Hormones are involved in a variety of different processes in the body and can vary greatly in their chemical composition and structure. The structures of some hormones are shown in

| Tab | le 21 of the Data Booklet. | |
|-----|--|-----|
| (a) | State the function of hormones. | [1] |
| | | |
| (b) | State the names of two functional groups present in both testosterone and progesterone. | [2] |
| | | |
| (c) | State the name of one functional group present in both estradiol and testosterone. | [1] |
| | | |
| (d) | Explain three different ways oral contraceptives taken by women prevent pregnancy. | [3] |
| | | |



Starch and cellulose are polysaccharides found in many plants.

(a) Compare the structures of starch and cellulose. [3]

Option C — Chemistry in industry and technology

(a)

(b)

C1. The Industrial Revolution was the result of large-scale extraction of iron from its ore and had significant impact worldwide.

while a mixture of limestone, coke and iron(III) oxide is introduced at the top.

In a blast furnace, a large volume of air is introduced under pressure near the bottom

(i) State the equation for the reaction of coke with air in the blast furnace. [1] (ii) The product formed in part (i) reacts with coke to produce carbon monoxide. Explain, giving an equation, why this reaction is important in the extraction of iron. Iron formed in the blast furnace contains about 5 % carbon, together with other impurities such as phosphorus, silicon and sulfur. Describe, using equations, the chemical principle behind the conversion of impure iron into steel using the basic oxygen converter and explain how the impurities are removed. [3]



| | polyethene (HDPE). Discuss two factors scientists would have considered in choosing these catalysts. | 1 |
|-----|--|---|
| | | |
| | | |
| | | |
| | | |
| | | |
| (b) | Describe a structural feature of low-density polyethene (LDPE) that explains why LDPE has a different melting point from that of HDPE. | L |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| (c) | State one environmental impact of the disposal of these polyethenes by using incineration. | |



C3. Fuel cells convert chemical energy directly into electrical energy that can be used in

applications ranging from spacecraft to remote weather stations.

(a) Describe the composition of the electrodes in a hydrogen-oxygen fuel cell. [1]

(b) State the half-equation at each electrode in the hydrogen-oxygen alkaline cell. [2]

Positive electrode (cathode):

Negative electrode (anode):

C4. It was over a hundred years after the accidental discovery of liquid crystals that liquid-crystal

displays (LCDs) came into common use in the 1990s. Liquid crystals are formed over a temperature range between the solid and the liquid state.

(a) Describe the nematic liquid-crystal phase in terms of the arrangement of the molecules. [2]

(b) Explain the effect of increasing the temperature on the nematic liquid crystal. [2]

Option D — Medicines and drugs

| | ichies and drugs are natural of synthetic substances used for their effects on the body. |
|-----|--|
| (a) | List two general effects of medicines and drugs on the functioning of the body. |
| | |
| | |
| | |
| (b) | Describe the placebo effect and state its importance in drug development. |



Medicines are classified by the effect each has on the body and the mind. One such example

is depressants. Describe one effect of depressants on the human body when a moderate dose is used and (a) one when a high dose is used. [2] Moderate dose: High dose: Breathalysers use acidified potassium dichromate(VI) to oxidize any ethanol vapour in (b) an individual's breath to ethanoic acid. (i) Deduce the half-equation for the oxidation of ethanol with water to form ethanoic acid and hydrogen ions. [1] Deduce the half-equation for the reduction of acidified dichromate(VI) ions to Cr³⁺(aq) and water. [1] State **two** other methods that can detect ethanol in the human body. (c) [1]



| (a) | Compare the structures of the two stimulants amphetamine and epinephrine (adrenaline). | |
|-----|--|---|
| | Similarity: | |
| | | |
| | | |
| | Difference: | |
| | | |
| | | |
| b) | Predict and explain which of these two stimulants would be more water soluble. | [|
| | | |
| | | |
| | | |
| | | |

(This question continues on the following page)




(Question D3 continued)

| Caff | feine and nicotine are also stimulants. | |
|-------|--|-----|
| (i) | Identify the type of amine that is present in both caffeine and nicotine. | [1] |
| | | |
| (ii) | Deduce whether an aqueous solution of caffeine is acidic, basic or neutral and explain your reasoning. | [2] |
| | | |
| | | |
| (iii) | Describe two effects of caffeine when it is consumed in large amounts. | [1] |
| | | |



- **D4.** Antibiotics treat infections by stopping the growth of bacteria or destroying them.
 - (a) Identify the side-chain by drawing a circle around the side-chain in the structure of benzyl penicillin given below (the structure of penicillin is given in Table 20 of the Data Booklet).

[1]



(b) Discuss **two** problems associated with the overprescription of penicillin and explain how these are overcome.

| | 3 | 7 |
|---|----|---|
| L | Τ. | J |

Option E — Environmental chemistry

- **E1.** Nitrogen dioxide and sulfur dioxide are two air pollutants.
 - Nitrogen dioxide is formed in a two-stage process. Describe one anthropogenic (man-made) source of nitrogen dioxide and state the two chemical equations for its formation. [2] Apart from using renewable technologies, state **one** method in each case for the removal of (b) nitrogen dioxide and sulfur dioxide from emission gases before they enter the atmosphere. [2] Nitrogen dioxide: Sulfur dioxide: Both of these air pollutants also contribute to acid deposition. State one chemical (c) equation for each gas to describe how each forms an acidic solution. [2]



The temperature of the Earth is increasing. There is considerable scientific evidence to

E2.

suggest this is due to an increase in the concentration of greenhouse gases as a result of human activity. Explain how this enhanced greenhouse effect causes the average temperature of the (a) Earth to increase. [3] (b) Compare the contributions of carbon dioxide and methane to the enhanced greenhouse effect. [2]

[4]

E3. In many countries the reserves of fresh water are scarce, so fresh water is obtained from sea water. Antigua, for example, produces much of its fresh water by reverse osmosis whereas the UAE produces much of its fresh water through multi-stage distillation.

Evaluate these two processes. Your answer should include at least **one** advantage and **one** disadvantage for each process.

| F | Advantage distillation: |
|---|-------------------------------|
| | |
| • | |
| Ι | Disadvantage distillation: |
| | |
| | |
| A | Advantage reverse osmosis: |
| | |
| | |
| Ι | Disadvantage reverse osmosis: |
| | |
| | |



[3]

E4. Ozone is a naturally occurring component of the stratosphere. The ozone layer occurs in the stratosphere about 30 km above the surface of the Earth.

Stratospheric ozone is in dynamic equilibrium with oxygen and is continually being formed and decomposed.

| (a) | State | the | condition | needed | for | the | depletion | of | ozone | in | the | stratosphere | by |
|-----|--------|--------|-------------|-------------------|------|------|--------------|-------|---------|-----|-----|--------------|----|
| | natura | ıl pro | cesses. Sta | ite the tw | o ch | emic | al equations | s for | the pro | ces | S. | | |

| |
|------|
| |
| |
| |
| |
| |

| (b) | State the names of two ozone-depleting pollutants and state their sources. | [2 |
|-----|---|----|
| | | |
| | | |
| | | |



Option F — Food chemistry

| (a) | State | e the empirical formula and structural features of monosaccharides. |
|-----|-------|---|
| | | |
| | | |
| (b) | (i) | State the structural formula of 2-aminoethanoic acid. |
| | (ii) | Deduce the structural formula of a triester formed from three long-chain carboxylic acid molecules, RCOOH, and one propane-1,2,3-triol molecule, HO-CH ₂ CH(OH)-CH ₂ OH. Identify one of the ester linkages in the structure by drawing a rectangle around it. |
| | | |
| | | |
| | | |

(This question continues on the following page)



Turn over

(Question F1 continued)

- (c) Examples of straight-chain fatty acids include $C_{19}H_{39}COOH$, $C_{19}H_{31}COOH$ and $C_{19}H_{29}COOH$.
 - (i) Deduce the number of C=C bonds present in one molecule of each fatty acid. [2]

| C ₁₉ H ₃₉ COOH: | |
|---------------------------------------|--|
| C ₁₉ H ₃₁ COOH: | |
| C ₁₉ H ₂₉ COOH: | |
| | |

| Deduce the least stable of the three fatty acids and explain your reasoning. |
|---|
| |
| |
| |
| |
| |
| |
| |

Food shelf life is the time it takes for a particular foodstuff to become unsuitable for eating

because it no longer meets customer or regulatory expectations. As a result, in many parts of

F2.

the world, packaged foods have a date before which they should be consumed. State the meaning of the term *rancidity* as it applies to fats. [1] (a) Rancidity in lipids occurs by hydrolytic and oxidative processes. (b) Compare the two rancidity processes. (i) [2] Hydrolytic process: Oxidative process:

(This question continues on the following page)



Turn over

(Question F2 continued)

| (ii) | State one example of a food containing lipids that undergoes hydrolytic rancidity and one that undergoes oxidative rancidity. | [2] |
|------|---|-----|
| | Hydrolytic rancidity: | |
| | | |
| | Oxidative rancidity: | |
| | | |
| | | |



F3. The browning of food during cooking is caused by two different processes, *caramelization* and *non-enzymatic browning* (Maillard reaction). These processes lead to a pleasant colour and flavour in a variety of food products. In the foods listed below, assume the following formulas:

• Sugar: $C_6H_{12}O_6$

• Reducing sugar: HOCH₂(CH(OH))₄CHO

• Amino acid: H₂N-CHR-COOH

Compare the two browning processes with respect to the chemical composition of the foods affected and state the chemical equation for the Maillard reaction.

[4]

| Chemical | composition of foods that undergo the caramelization process: |
|------------|---|
| | |
| | |
| | |
| Chemical | composition of foods that undergo the non-enzymatic browning process: |
| | |
| | |
| | |
| Maillard r | reaction equation: |
| | |
| | |
| | |
| | |
| | |



Foods derived from genetically modified organisms were introduced in the early 1990s.

F4.

State **one** benefit and **one** concern of consuming genetically modified foods.

[2]

Benefit:


Concern:



Option G — Further organic chemistry

G1. Consider the following reaction pathway starting with compound **A** which is but-1-ene.

(a) (i) A reacts with HBr to form a major and a minor product. Draw the structural formulas of these two isomers.



(ii) Explain the mechanism of the reaction of **A** with HBr to form the major product, **B**, using curly arrows to represent the movement of electron pairs. [4]

(This question continues on the following page)



Turn over

| (Ouestion | GI | continued, |
|-----------|----|------------|
| | | |

| | | | | | | | | | | | |
|---------|---------|---------|---------|--------|-------|---------------|--------|-------|-------|---------|------|
| | | | | | | | | | | | |
| | | | | | | | | | | | |
| | | | | | | | | | | | |
| | | | | | | | | | | | |
| | | | | | | | | | | | |
| | | | | | | | | | | | |
| pound C | e conve | erted d | irectly | to com | pound | A . Io | dentif | y the | reage | ent use | d fo |
| pound C | e conve | erted d | irectly | to com | pound | A . Io | dentif | y the | reage | ent use | d fo |
| | e conve | erted d | irectly | to com | pound | A . Io | dentif | y the | reage | ent use | d fo |
| | e conve | erted d | irectly | to com | pound | A . Io | dentif | y the | reage | ent use | d fo |

(This question continues on the following page)



(Question G1 continued)

| (i) | Deduce the structural formula of D . | |
|-------|---|---|
| | | |
| | | |
| | | |
| | | |
| | | |
| | | _ |
| (ii) | Compound D reacts with 2,4-dinitrophenylhydrazine, $H_2NNH-C_6H_3(NO_2)_2$, to give | |
| | a solid E . Deduce the structural formula of E . | |
| | | |
| | | |
| | | |
| | | |
| | | |
| (iii) | Explain the mechanism for the reaction of D with hydrogen cyanide to form F , | |
| () | using curly arrows to represent the movement of electron pairs. | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |



| ene, C_6H_6 , is the simplest aromatic hydrocarbon. ribe and explain the structure of benzene. | [3] |
|--|-----|
| ene, C_6H_6 , is the simplest aromatic hydrocarbon. | [3] |
| ene, C_6H_6 , is the simplest aromatic hydrocarbon. | [3] |
| | [3] |
| | [3] |
| | [3] |
| | [3] |
| | [3] |
| | [3] |
| | [3] |
| | [3] |
| 1 | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |

