



MARKSCHEME

May 2007

MATHEMATICS

Higher Level

Paper 1

*This markscheme is **confidential** and for the exclusive use of examiners in this examination session.*

*It is the property of the International Baccalaureate and must **not** be reproduced or distributed to any other person without the authorization of IBCA.*

Instructions to Examiners

Abbreviations

- M** Marks awarded for attempting to use a correct **Method**; working must be seen.
- (M)** Marks awarded for **Method**; may be implied by **correct** subsequent working.
- A** Marks awarded for an **Answer** or for **Accuracy**: often dependent on preceding **M** marks.
- (A)** Marks awarded for an **Answer** or for **Accuracy**; may be implied by **correct** subsequent working.
- R** Marks awarded for clear **Reasoning**.
- N** Marks awarded for **correct** answers if **no** working shown.
- AG** Answer given in the question and so no marks are awarded.

Using the markscheme

1 General

Write the marks in red on candidates' scripts, in the right hand margin.

- Show the **breakdown** of individual marks awarded using the abbreviations **MI**, **AI**, etc.
- Write down the total for each **question** (at the end of the question) and **circle** it.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is not possible to award **M0** followed by **AI**, as **A** mark(s) depend on the preceding **M** mark(s), if any.
- Where **M** and **A** marks are noted on the same line, e.g. **MIAI**, this usually means **MI** for an **attempt** to use an appropriate method (e.g. substitution into a formula) and **AI** for using the **correct** values.
- Where the markscheme specifies (**M2**), **N3**, etc., do **not** split the marks.
- Once a correct answer to a question or part-question is seen, ignore further working.

3 N marks

*Award N marks for **correct** answers where there is **no** working.*

- Do **not** award a mixture of **N** and other marks.
- There may be fewer **N** marks available than the total of **M**, **A** and **R** marks; this is deliberate as it penalizes candidates for not following the instruction to show their working.

4 Implied marks

*Implied marks appear in **brackets e.g. (MI)**, and can only be awarded if **correct** work is seen or if implied in subsequent working.*

- Normally the correct work is seen or implied in the next line.
- Marks **without** brackets can only be awarded for work that is **seen**.

5 Follow through marks

*Follow through (**FT**) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s). To award **FT** marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part.*

- If the question becomes much simpler because of an error then use discretion to award fewer **FT** marks.
- If the error leads to an inappropriate value (*e.g.* $\sin\theta=1.5$), do not award the mark(s) for the final answer(s).
- Within a question part, once an error is made, no further **dependent A** marks can be awarded, but **M** marks may be awarded if appropriate.
- Exceptions to this rule will be explicitly noted on the markscheme.

6 Mis-read

*If a candidate incorrectly copies information from the question, this is a mis-read (**MR**). Apply a **MR** penalty of 1 mark to that question. Award the marks as usual and then write $-1(\mathbf{MR})$ next to the total. Subtract 1 mark from the total for the question. A candidate should be penalized only once for a particular mis-read.*

- If the question becomes much simpler because of the **MR**, then use discretion to award fewer marks.
- If the **MR** leads to an inappropriate value (*e.g.* $\sin\theta=1.5$), do not award the mark(s) for the final answer(s).

7 Discretionary marks (*d*)

*An examiner uses discretion to award a mark on the rare occasions when the markscheme does not cover the work seen. The mark should be labelled (**d**) and a brief **note** written next to the mark explaining this decision.*

8 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If in doubt, contact your team leader for advice.

- Alternative methods for complete questions are indicated by **METHOD 1, METHOD 2, etc.**
- Alternative solutions for part-questions are indicated by **EITHER . . . OR.**
- Where possible, alignment will also be used to assist examiners in identifying where these alternatives start and finish.

9 Alternative forms

Unless the question specifies otherwise, **accept** equivalent forms.

- As this is an international examination, accept all alternative forms of **notation**.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, **simplified** answers, (which candidates often do not write in examinations), will generally appear in brackets. Marks should be awarded for either the form preceding the bracket or the form in brackets (if it is seen).

Example: for differentiating $f(x) = 2\sin(5x - 3)$, the markscheme gives:

$$f'(x) = 2\cos(5x - 3) \cdot 5 = 10\cos(5x - 3) \quad \mathbf{AI}$$

Award **AI** for $2\cos(5x - 3) \cdot 5$, even if $10\cos(5x - 3)$ is not seen.

10 Accuracy of Answers

If the level of accuracy is specified in the question, a mark will be allocated for giving the answer to the required accuracy.

- **Rounding errors:** only applies to final answers not to intermediate steps.
- **Level of accuracy:** when this is not specified in the question the general rule applies: *unless otherwise stated in the question all numerical answers must be given exactly or correct to three significant figures.*

Candidates should be penalized **once only IN THE PAPER** for an accuracy error (**AP**). Award the marks as usual then write (**AP**) against the answer. On the **front** cover write $-1(\mathbf{AP})$. Deduct 1 mark from the total for the paper, not the question.

- If a final correct answer is incorrectly rounded, apply the **AP**.
- If the level of accuracy is not specified in the question, apply the **AP** for correct answers not given to three significant figures.

If there is no working shown, and answers are given to the correct two significant figures, apply the **AP**. However, do **not** accept answers to one significant figure without working.

11 Crossed out work

If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work.

QUESTION 1

METHOD 1

$\hat{BAC} = 80^\circ$	<i>(AI)</i>	
Using the sine rule to find AB	<i>MI</i>	
$\frac{105}{\sin 80^\circ} = \frac{AB}{\sin 40^\circ}$	<i>AIAI</i>	
Attempting to solve for AB	<i>(MI)</i>	
AB = 68.5 (m)	<i>AI</i>	<i>N4</i>

METHOD 2

$\hat{BAC} = 80^\circ$	<i>(AI)</i>	
Using the sine rule to find AC	<i>MI</i>	
$\frac{105}{\sin 80^\circ} = \frac{AC}{\sin 60^\circ}$	<i>AI</i>	
$AC = \frac{105 \sin 60^\circ}{\sin 80^\circ} \quad (=92.335\dots)$	<i>AI</i>	
Using the cosine rule to find AB <i>i.e.</i> $AB^2 = 105^2 + AC^2 - 2(105)(AC)\cos 40^\circ$	<i>(MI)</i>	
AB = 68.5 (m)	<i>AI</i>	<i>N4</i>

QUESTION 2

(a) Recognising how to find the median (25.5 th item)	<i>(MI)</i>	
median = $\frac{2+3}{2}$	<i>(AI)</i>	
$= 2.5$	<i>AI</i>	<i>N3</i>
(b) Using $\bar{x} = \frac{\sum xf}{n}$	<i>(MI)</i>	
$\bar{x} = \frac{(0 \times 7) + (1 \times 3) + (2 \times 15) + (3 \times 11) + (4 \times 6) + (5 \times 5) + (6 \times 3)}{50}$	<i>(AI)</i>	
$\bar{x} = \frac{133}{50} = 2.66$	<i>AI</i>	<i>N3</i>

QUESTION 3

$$\mathbf{n}_1 = -2\mathbf{i} + \mathbf{j} - \mathbf{k} \text{ and } \mathbf{n}_2 = \mathbf{i} + 2\mathbf{j} - \mathbf{k}$$

(AI)(AI)

$$|\mathbf{n}_1| = \sqrt{6} \text{ and } |\mathbf{n}_2| = \sqrt{6}$$

(AI)

$$\cos\theta = \frac{\mathbf{n}_1 \cdot \mathbf{n}_2}{|\mathbf{n}_1| |\mathbf{n}_2|}$$

$$\cos\theta = \frac{(-2\mathbf{i} + \mathbf{j} - \mathbf{k}) \cdot (\mathbf{i} + 2\mathbf{j} - \mathbf{k})}{\sqrt{6} \times \sqrt{6}}$$

MI(AI)

$$\cos\theta = \frac{1}{6} \text{ (0.167 to 3 s.f.)}$$

AI

N4

QUESTION 4

$$2(\ln x)^2 - 3\ln x + 1 = 0$$

(AI)

Attempting to factorise or using the quadratic formula

(MI)

$$\ln x = \frac{1}{2}, \ln x = 1$$

AIAI

$$x = \sqrt{e}, x = e$$

AIAI

N2

QUESTION 5

Separating variables

(MI)

$$\int \frac{dy}{y^2} = \int 2x \, dx$$

AI

$$-\frac{1}{y} = x^2 + C$$

AIAI

Note: The first AI above is for a correct LHS and the second AI is for a correct RHS that must include C.

Using $y(0) = 1$ gives $C = -1 \left(-\frac{1}{y} = x^2 - 1 \right)$

MI

$$y = -\frac{1}{x^2 - 1} \left(= \frac{1}{1 - x^2} \right)$$

AI

N0

QUESTION 6**METHOD 1**

$$\vec{AB} = 2\mathbf{j} - \mathbf{k} \text{ and } \vec{AC} = -3\mathbf{i} + 2\mathbf{j} \quad (AI)(AI)$$

$$\vec{AB} \times \vec{AC} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 2 & -1 \\ -3 & 2 & 0 \end{vmatrix} \quad MI$$

$$= 2\mathbf{i} + 3\mathbf{j} + 6\mathbf{k} \quad AI$$

$$\text{Area } \Delta ABC = \frac{1}{2} \left| \vec{AB} \times \vec{AC} \right| \quad (MI)$$

$$= \frac{7}{2} \quad AI \quad N4$$

METHOD 2

$$\vec{AB} = 2\mathbf{j} - \mathbf{k} \text{ and } \vec{AC} = -3\mathbf{i} + 2\mathbf{j} \quad (AI)(AI)$$

$$\text{Attempting to use the scalar product to find } \theta \text{ i.e. } \vec{AB} \cdot \vec{AC} = \left| \vec{AB} \right| \left| \vec{AC} \right| \cos \theta \quad MI$$

$$\cos \theta = \frac{4}{\sqrt{65}} \left(\theta = \arccos \frac{4}{\sqrt{65}} = \arcsin \frac{7}{\sqrt{65}} = 60.255\dots^\circ \right) \quad AI$$

$$\text{Area } \Delta ABC = \frac{1}{2} \left| \vec{AB} \right| \left| \vec{AC} \right| \sin \theta \left(= \frac{1}{2} \times \sqrt{5} \times \sqrt{13} \times \frac{7}{\sqrt{65}} \right) \quad (MI)$$

$$= \frac{7}{2} \quad AI \quad N4$$

QUESTION 7

$$(a) \quad X \sim B\left(7, \frac{1}{5}\right) \quad (AI)$$

$$P(X = 4) = \binom{7}{4} \times \left(\frac{1}{5}\right)^4 \times \left(\frac{4}{5}\right)^3 \quad (MI)$$

$$= 0.0287 \quad \left(= \frac{448}{15625} \right) \quad AI \quad N3$$

$$(b) \quad P(\text{pass}) = P(X \geq 4) \text{ OR } 1 - P(X \leq 3) \text{ (or equivalent)} \quad A2$$

$$= 0.0333 \quad \left(= \frac{521}{15625} \right) \quad AI \quad N2$$

Note: Accept 0.0334

QUESTION 8

(a) Attempting to find $\det A$ **(MI)**
 $\det A = k^2 + 2k - 1$ **AI** **N2**

(b) **METHOD 1**

Solving $k^2 + 2k - 1 = 0$ or equivalent for k **MI**
 $k = -1 \pm \sqrt{2}$ $(-2.41, 0.414)$ **(AI)**
 System has a unique solution provided $\det A \neq 0$ **(RI)**
 $k \in \mathbb{R} \setminus \{-1 \pm \sqrt{2}\}$ (accept $k \neq -1 \pm \sqrt{2}, k \neq -2.41, 0.414$) **AI** **N3**

METHOD 2

System has a unique solution provided $\det A \neq 0$ **(RI)**
 $k^2 + 2k - 1 \neq 0$ **(AI)**
 Solving $k^2 + 2k - 1 \neq 0$ or equivalent for k **MI**
 $k \in \mathbb{R} \setminus \{-1 \pm \sqrt{2}\}$ (accept $k \neq -1 \pm \sqrt{2}, k \neq -2.41, 0.414$) **AI** **N3**

QUESTION 9

METHOD 1

$s = \int (3t^2 - 4t + 2) dt$ **AI**
 Attempting to integrate the RHS **(MI)**
 $s = t^3 - 2t^2 + 2t + C$ **AI**

Note: The **AI** above must include C .

Using $s(0) = -3$ gives $C = -3$ ($s = t^3 - 2t^2 + 2t - 3$) **AI**
 Solving $t^3 - 2t^2 + 2t - 3 = 0$ for t **(MI)**
 $t = 1.81$ (sec) **AI** **N3**

METHOD 2

$\int_0^T (3t^2 - 4t + 2) dt = \int_{-3}^0 ds$ ($=3$) **AIAI**
 Solving for T **(MI)**
 $T = 1.81$ (sec) **A3** **N3**

QUESTION 10

(a) Mode = 0

AI

(b) $E(X) = \frac{8}{\pi} \int_0^2 \frac{x}{x^2 + 4} dx$

MIAI

Attempting substitution method or using $\int \frac{f'(x)}{f(x)} dx = \ln f(x)$

(MI)

$E(X) = \frac{4}{\pi} \ln u \Big|_4^8$ or $E(X) = \frac{4}{\pi} [\ln(x^2 + 4)]_0^2$

(AI)

$= \frac{4}{\pi} (\ln 8 - \ln 4) \left(= \frac{4}{\pi} \ln \left(\frac{8}{4} \right), \frac{4}{\pi} \ln 2 \right)$

AI

N0

QUESTION 11

METHOD 1

Using factor theorem

(MI)

Substituting $z = -1 - i$ into $P(z)$

MI

$-(6 + n) + (2m - 2 - n)i = 0$

AI

Equating both real and imaginary parts to zero

MI

Hence $m = -2$ and $n = -6$

AIAI

N2

METHOD 2

Using Conjugate root theorem

MI

Multiply $(z + 1 - i)(z + 1 + i) = z^2 + 2z + 2$

MI

Let $P(z) = (z^2 + 2z + 2)(z - a)$

(MI)

$-2a = -8 \quad a = 4$

AI

Hence $m = -2$ and $n = -6$

AIAI

N2

QUESTION 12

METHOD 1

$$\Delta = 4 - 4k(3k + 2) \quad (= -12k^2 - 8k + 4, = -4(k + 1)(3k - 1))$$

MIAI

$$\Delta = 0 \Rightarrow k = -1, k = \frac{1}{3}$$

(AI)

For 2 distinct roots, $\Delta > 0$

(RI)

$$-1 < k < \frac{1}{3}$$

A2

N4

METHOD 2

For 2 distinct roots, $\Delta > 0$

(RI)

$$\Delta = 4 - 4k(3k + 2) \quad (= -12k^2 - 8k + 4, = -4(k + 1)(3k - 1))$$

MIAI

$$\Delta = 0 \Rightarrow k = -1, k = \frac{1}{3}$$

(AI)

$$-1 < k < \frac{1}{3}$$

A2

N4

QUESTION 13

Using $S_n = \frac{n}{2} 2u_1 + (n-1)d$ with $u_1 = -6$ and $d = 7$

(MI)(AI)

$$S_n = \frac{n}{2}(7n - 19) \text{ or equivalent}$$

AI

Solving $S_n > 10\,000$ or equivalent for n for *e.g.* $(7n^2 - 19n - 20\,000 > 0)$
 $n > 54.8\dots$

MI

(AI)

The least number of terms is 55.

AI

N4

QUESTION 14

- Recognition of stretch or compression parallel to x -axis *MI*
- Scale factor is $\frac{6}{\pi}$ or $\frac{\pi}{6}$ respectively *AI*
- Reflection in x -axis *AI*
- Recognition of stretch parallel to y -axis with scale factor 2 *AI*
- Recognition of translation $\begin{pmatrix} 0 \\ 8 \end{pmatrix}$ *AI*
- A correct sequence (i.e. the translation must be stated last). *AI*


QUESTION 15

- $V = \pi \int_a^b y^2 dx$ *(MI)*
- $V = \pi \int_0^{\frac{\pi}{4}} \sin^2 3x dx$ *AI*
- $\sin^2 3x = \frac{1}{2}(1 - \cos 6x)$ *(AI)*
- $V = \frac{\pi}{2} \int_0^{\frac{\pi}{4}} (1 - \cos 6x) dx$
- $= \frac{\pi}{2} \left[x - \frac{1}{6} \sin 6x \right]_0^{\frac{\pi}{4}}$ *AI*
- $= \frac{\pi}{2} \left(\frac{\pi}{4} + \frac{1}{6} \right) \left(= \frac{\pi^2}{8} + \frac{\pi}{12} \right)$ *MI AI* *N0*

QUESTION 16

- (a) $P(X > 45) = P(Z < 1.25)$ *(MI)*
- $= 0.894$ *AI* *N2*
- (b) Using conditional probability $P(X > 55 | X > 45)$ *(AI)*
- $= \frac{P(X > 55 \cap X > 45)}{P(X > 45)}$ *(AI)*
- $= \frac{P(X > 55)}{P(X > 45)} \left(= \frac{0.1056...}{0.8943...} \right)$ *(AI)*
- FT** numerator = 1 – (their answer to a).
- $= 0.118$ *A1* *N2*

QUESTION 17



AIAI

(b) $A = \int_a^b g(x) - f(x) \, dx$ (using an appropriate definite integral)

(AI)

$a = 0.50546\dots, b = 1.227\dots$

(AI)(AI)

$A = 0.201$

AI

N2

QUESTION 18

(a) Use of quotient (or product) rule

(MI)

$$f'(x) = \frac{2(x^2 + 6) - (2x \times 2x)}{(x^2 + 6)^2} = \frac{2x(-1)(x^2 + 6)^{-2}(2x) + 2(x^2 + 6)^{-1}}{(x^2 + 6)^2}$$

AI

$$= \frac{12 - 2x^2}{(x^2 + 6)^2}$$

AG

N0

(b) Solving $f'(x) = 0$ for x

(MI)

$x = \pm\sqrt{6}$

AI

f has to be 1-1 for f^{-1} to exist and so the least value of b is the larger of the two x -coordinates (accept a labelled sketch)

RI

Hence $b = \sqrt{6}$

AI

N2

QUESTION 19

Let $u = e^x$

MI

$du = e^x dx$ (or equivalent)

AI

When $x = 0$, $u = 1$ **and** when $x = \ln 3$, $u = 3$

(AI)

$$\int_0^{\ln 3} \frac{e^x}{e^{2x} + 9} dx = \int_1^3 \frac{1}{u^2 + 9} du$$

AI

$$= \frac{1}{3} \left[\arctan \frac{u}{3} \right]_1^3$$

AI

$$= \frac{1}{3} \left(\arctan 1 - \arctan \frac{1}{3} \right) \quad \left(= \frac{\pi}{12} - \frac{1}{3} \arctan \frac{1}{3}, \frac{1}{3} \arctan \frac{1}{2} \right)$$

AI

NO

QUESTION 20

$z^2 = x^2 + y^2$ (or equivalent)

(MI)

$z = \sqrt{0.8^2 + 0.6^2}$ (=1, initially)

AI

Attempting to differentiate implicitly with respect to t

MI

$$2z \frac{dz}{dt} = 2x \frac{dx}{dt} + 2y \frac{dy}{dt}$$

AI

$$\frac{dz}{dt} = -(0.8 \times 60) - (0.6 \times 70)$$

AI

Rate is $-90 \text{ (km h}^{-1}\text{)}$

AI

NO