



MARKSCHEME

May 2013

**MATHEMATICS
DISCRETE MATHEMATICS**

Higher Level

Paper 3

*This markscheme is **confidential** and for the exclusive use of
examiners in this examination session.*

*It is the property of the International Baccalaureate and
must **not** be reproduced or distributed to any other person
without the authorization of the IB Assessment Centre.*

1. (a) using the Euclidean Algorithm,

$$332 = 3 \times 99 + 35$$

$$99 = 2 \times 35 + 29$$

$$35 = 1 \times 29 + 6$$

$$29 = 4 \times 6 + 5$$

$$6 = 1 \times 5 + 1$$

hence 332 and 99 have a gcd of 1

MI

A1

A1

A1

AG

Note: For both (a) and (b) accept layout in tabular form, especially the brackets method of keeping track of the linear combinations as the method proceeds.

[4 marks]

- (b) (i) working backwards,

$$6 - 5 = 1$$

$$6 - (29 - 4 \times 6) = 1 \text{ or } 5 \times 6 - 29 = 1$$

$$5 \times (35 - 29) - 29 = 1 \text{ or } 5 \times 35 - 6 \times 29 = 1$$

$$5 \times 35 - 6 \times (99 - 2 \times 35) = 1 \text{ or } 17 \times 35 - 6 \times 99 = 1$$

$$17 \times (332 - 3 \times 99) - 6 \times 99 = 1 \text{ or } 17 \times 332 - 57 \times 99 = 1$$

a solution to the Diophantine equation is therefore

$$x = 17, y = 57$$

the general solution is

$$x = 17 + 99N, y = 57 + 332N$$

(MI)

A1

A1

A1

(A1)

A1A1

Note: If part (a) is wrong it is inappropriate to give *FT* in (b) as the numbers will contradict, however the *MI* can be given.

- (ii) it follows from previous work that

$$17 \times 332 = 1 + 99 \times 57$$

$$\equiv 1 \pmod{57}$$

$z = 332$ is a solution to the given congruence

the general solution is $332 + 57N$ so the smallest solution is 47

(MI)

(A1)

(A1)

A1

A1

[11 marks]

Total [15 marks]

2. (a) (i) there is an Eulerian trail because there are only 2 vertices of odd degree ***R1***
 there is no Eulerian circuit because not all vertices have even degree ***R1***

(ii) eg GBAGFBCFECDE ***A2***

[4 marks]

(b)	(i)	Step	Vertices labelled	Working values	
		1	A	A(0), B-3, G-2	<i>M1A1</i>
		2	A, G	A(0), G(2), B-3, F-8	<i>A1</i>
		3	A, G, B	A(0), G(2), B(3), F-7, C-10	<i>A1</i>
		4	A, G, B, F	A(0), G(2), B(3), F(7), C-9, E-12	
		5	A, G, B, F, C	A(0), G(2), B(3), F(7), C(9), E-10, D-15	<i>A1</i>
		6	A, G, B, F, C, E	A(0), G(2), B(3), F(7), C(9), E(10), D-14	
		7	A, G, B, F, C, E, D	A(0), G(2), B(3), F(7), C(9), E(10), D(14)	<i>A1</i>

Note: In both (i) and (ii) accept the tabular method including back tracking or labels by the vertices on a graph.

Note: Award ***M1A1A1A1A0A0*** if final labels are correct but intermediate ones are not shown.

(ii) minimum weight path is ABFCED ***A1***
 minimum weight is 14 ***A1***

Note: Award the final two ***A1*** marks whether or not Dijkstra's Algorithm is used.

[8 marks]

Total [12 marks]

3. (a) the equation can be written as

$$(3n+3)^2 = n^3 + 3n^2 + 3n + 1$$

any valid method of solution giving $n=8$

M1A1

(M1)A1

Note: Attempt to change at least one side into an equation in n gains the ***M1***.

[4 marks]

(b) **METHOD 1**

as decimal numbers,

$$(33)_8 = 27, (1331)_8 = 729$$

converting to base 7 numbers,

$$\begin{array}{r} 27 = (36)_7 \\ 7) \underline{729} \\ 7) \underline{104}(1 \\ 7) \underline{14}(6 \\ 7) \underline{2}(0 \\ 7) \underline{0}(2 \end{array}$$

$$\text{therefore } 729 = (2061)_7$$

the required equation is

$$36^2 = 2061$$

A1A1

A1

MI

A1

A1

METHOD 2

as a decimal number, $(33)_8 = 27$

A1

converting to base 7,

$$27 = (36)_7$$

multiplying base 7 numbers

$$\begin{array}{r} 36 \\ \times 36 \\ \hline 1440 \\ 321 \\ \hline 2061 \end{array}$$

the required equation is

$$36^2 = 2061$$

A1

MI

A1

A1

A1

Note: Allow ***M1*** for showing the method of converting a number to base 7 regardless of what number they convert.

[6 marks]

Total [10 marks]

4. (a) evaluating the adjacency matrix to the fifth power
number of walks = 14

(M1)
A2

[3 marks]

- (b) number of edges in $G = 5$

$$\text{number of edges in } G' = \binom{5}{2} - 5 \\ = 5$$

A1
(M1)
A1

Note: Allow listing of edges in G' or drawing graphs.

[3 marks]

- (c) (i) the adjacency matrix of G' is

	B	D	A	C	E
B	0	1	0	1	1
D	1	0	0	0	0
A	0	0	0	1	0
C	1	0	1	0	1
E	1	0	0	1	0

A4

Note: Award A3 for one error, A2 for two errors, A1 for three errors and A0 for more than three errors.

- (ii) it follows that G and G' are isomorphic because the adjacency matrices of G and G' are identical

R1

Note: Or equivalent comprehensive explanation.

[5 marks]

- (d) let H have e edges

M1

$$\text{number of edges in } H' = \binom{6}{2} - e = 15 - e$$

A1

for an isomorphism to exist, these must be equal:

M1

$$e = 15 - e \Rightarrow e = 7.5$$

A1

which is impossible so no isomorphism

AG

[4 marks]

Total [15 marks]

5. (a) using Fermat's little theorem,

$$k^p \equiv k \pmod{p} \quad (\text{M1})$$

therefore,

$$\sum_{k=1}^p k^p \equiv \sum_{k=1}^p k \pmod{p} \quad \text{M1}$$

$$\equiv \frac{p(p+1)}{2} \pmod{p} \quad \text{A1}$$

$$\equiv 0 \pmod{p} \quad \text{AG}$$

since $\frac{(p+1)}{2}$ is an integer (so that the right-hand side is a multiple of p) **R1**

[4 marks]

- (b) using the alternative form of Fermat's little theorem,

$$k^{p-1} \equiv 1 \pmod{p}, 1 \leq k \leq p-1 \quad \text{A1}$$

$$k^{p-1} \equiv 0 \pmod{p}, k = p \quad \text{A1}$$

therefore,

$$\sum_{k=1}^p k^{p-1} \equiv \sum_{k=1}^{p-1} 1 (+0) \pmod{p} \quad \text{M1}$$

$$\equiv p-1 \pmod{p} \quad \text{A1}$$

(so $n = p-1$)

Note: Allow first **A1** even if qualification on k is not given.

[4 marks]

Total [8 marks]