



MARKSCHEME

November 2013

MATHEMATICS SETS, RELATIONS AND GROUPS

Higher Level

Paper 3

*This markscheme is **confidential** and for the exclusive use of examiners in this examination session.*

*It is the property of the International Baccalaureate and must **not** be reproduced or distributed to any other person without the authorization of the IB Assessment Centre.*

Instructions to Examiners

Abbreviations

- M** Marks awarded for attempting to use a correct **Method**; working must be seen.
- (M)** Marks awarded for **Method**; may be implied by **correct** subsequent working.
- A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- (A)** Marks awarded for an **Answer** or for **Accuracy**; may be implied by **correct** subsequent working.
- R** Marks awarded for clear **Reasoning**.
- N** Marks awarded for **correct** answers if **no** working shown.
- AG** Answer given in the question and so no marks are awarded.

Using the markscheme

1 General

Mark according to Scoris instructions and the document “**Mathematics HL: Guidance for e-marking November 2013**”. It is **essential** that you read this document before you start marking. In particular, please note the following.

Marks must be recorded using the annotation stamps. Please check that you are entering marks for the right question.

- If a part is **completely correct**, (and gains all the “must be seen” marks), use the ticks with numbers to stamp full marks.
- If a part is completely wrong, stamp **A0** by the final answer.
- If a part gains anything else, it **must** be recorded using **all** the annotations.

All the marks will be added and recorded by Scoris.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is not possible to award **M0** followed by **AI**, as **A** mark(s) depend on the preceding **M** mark(s), if any.
- Where **M** and **A** marks are noted on the same line, for example, **MIAI**, this usually means **MI** for an **attempt** to use an appropriate method (for example, substitution into a formula) and **AI** for using the **correct** values.
- Where the markscheme specifies (**M2**), **N3**, etc, do **not** split the marks.
- Once a correct answer to a question or part-question is seen, ignore further working.

3 ***N* marks**

Award ***N*** marks for **correct** answers where there is **no** working.

- Do **not** award a mixture of ***N*** and other marks.
- There may be fewer ***N*** marks available than the total of ***M***, ***A*** and ***R*** marks; this is deliberate as it penalizes candidates for not following the instruction to show their working.

4 **Implied marks**

Implied marks appear in brackets, for example, (MI), and can only be awarded if correct work is seen or if implied in subsequent working.

- Normally the correct work is seen or implied in the next line.
- Marks **without** brackets can only be awarded for work that is **seen**.

5 **Follow through marks**

Follow through (FT) marks are awarded where an incorrect answer from one part of a question is used correctly in subsequent part(s). To award FT marks, there must be working present and not just a final answer based on an incorrect answer to a previous part.

- If the question becomes much simpler because of an error then use discretion to award fewer **FT** marks.
- If the error leads to an inappropriate value (for example, $\sin\theta = 1.5$), do not award the mark(s) for the final answer(s).
- Within a question part, once an error is made, no further **dependent A** marks can be awarded, but **M** marks may be awarded if appropriate.
- Exceptions to this rule will be explicitly noted on the markscheme.

6 **Mis-read**

If a candidate incorrectly copies information from the question, this is a mis-read (MR). A candidate should be penalized only once for a particular mis-read. Use the MR stamp to indicate that this has been a mis-read. Then deduct the first of the marks to be awarded, even if this is an M mark, but award all others so that the candidate only loses one mark.

- If the question becomes much simpler because of the **MR**, then use discretion to award fewer marks.
- If the **MR** leads to an inappropriate value (for example, $\sin\theta = 1.5$), do not award the mark(s) for the final answer(s).

7 **Discretionary marks (d)**

An examiner uses discretion to award a mark on the rare occasions when the markscheme does not cover the work seen. In such cases the annotation DM should be used and a brief note written next to the mark explaining this decision.

8 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If in doubt, contact your team leader for advice.

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, etc.
- Alternative solutions for part-questions are indicated by **EITHER . . . OR**.
- Where possible, alignment will also be used to assist examiners in identifying where these alternatives start and finish.

9 Alternative forms

Unless the question specifies otherwise, **accept** equivalent forms.

- As this is an international examination, accept all alternative forms of **notation**.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, **simplified** answers, (which candidates often do not write in examinations), will generally appear in brackets. Marks should be awarded for either the form preceding the bracket or the form in brackets (if it is seen).

Example: for differentiating $f(x) = 2\sin(5x - 3)$, the markscheme gives:

$$f'(x) = (2\cos(5x - 3))5 \quad (=10\cos(5x - 3)) \quad \text{AI}$$

Award **AI** for $(2\cos(5x - 3))5$, even if $10\cos(5x - 3)$ is not seen.

10 Accuracy of Answers

Candidates should **NO LONGER** be penalized for an accuracy error (**AP**).

If the level of accuracy is specified in the question, a mark will be allocated for giving the answer to the required accuracy. When this is not specified in the question, all numerical answers should be given exactly or correct to three significant figures. Please check work carefully for **FT**.

11 Crossed out work

If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work.

12 Calculators

A GDC is required for paper 3, but calculators with symbolic manipulation features (for example, TI-89) are not allowed.

Calculator notation

The Mathematics HL guide says:

Students must always use correct mathematical notation, not calculator notation.

Do **not** accept final answers written using calculator notation. However, do not penalize the use of calculator notation in the working.


13 More than one solution

Where a candidate offers two or more different answers to the same question, an examiner should only mark the first response unless the candidate indicates otherwise.

1. (a) **Method 1**

sketch of the graph of f

(M1)



range of $f =$ co-domain, therefore f is surjective

RI

graph of f passes the horizontal line test, therefore f is injective

RI

therefore f is bijective

AG

[3 marks]

Note: Other explanations may be given (eg use of derivative or description of parabola).

Method 2

Injective: $f(a) = f(b) \Rightarrow a = b$

MI

$$(a-1)(a+2) = (b-1)(b+2)$$

$$a^2 + a = b^2 + b$$

solving for a by completing the square, or the quadratic formula,
 $a=b$

AI

surjective: for all $y \in \mathbb{R}^+$ there exists $x \in]1, \infty[$ such that $f(x) = y$

solving $y = x^2 + x - 2$ for x , $x = \frac{\sqrt{4y+9}-1}{2}$. For all positive real y , the

minimum value for $\sqrt{4y+9}$ is 3. Hence, $x \geq 1$

RI

since f is both injective and surjective, f is bijective.

AG

Method 3

f is bijective if and only if f has an inverse **(M1)**

solving $y = x^2 + x - 2$ for x , $x = \frac{\sqrt{4y+9}-1}{2}$. For all positive real y , the minimum value for $\sqrt{4y+9}$ is 3. Hence, $x \geq 1$ **RI**

$$f^{-1}(x) = \frac{\sqrt{4x+9}-1}{2} \quad \text{RI}$$

f has an inverse, hence f is bijective **AG**

[3 marks]

(b) (i) attempt to find counterexample **(M1)**
 eg $g(x, y) = g(y, x)$, $x \neq y$ **AI**

g is not injective **RI**

(ii) $-1 \leq \sin(x+y) \leq 1$ **(M1)**
 range of g is $[-1, 1] \times \mathbb{R} \neq \mathbb{R} \times \mathbb{R}$ **AI**
 g is not surjective **RI**

[6 marks]

continued ...

Question 1 continued

(c) let $h(x, y) = (u, v)$

then $u = x + 3y$

$v = 2x + y$

(M1)

solving simultaneous equations

(M1)

$$\text{eg } \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 & 3 \\ 2 & 1 \end{pmatrix}^{-1} \begin{pmatrix} u \\ v \end{pmatrix} \Rightarrow \begin{pmatrix} x \\ y \end{pmatrix} = -\frac{1}{5} \begin{pmatrix} 1 & -3 \\ -2 & 1 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix}$$

$$x = \frac{-u + 3v}{5} \quad y = \frac{2u - v}{5}$$

AI

$$\text{hence } h^{-1}(x, y) = \left(\frac{-x + 3y}{5}, \frac{2x - y}{5} \right)$$

AI

as this expression is defined for any values of $(x, y) \in \mathbb{R} \times \mathbb{R}$

RI

the inverse of h exists

AG

[5 marks]

Total [14 marks]

2. (a) (i) the order of a is a divisor of the order of G **(M1)**
 since the order of G is 12, the order of a must be 1, 2, 3, 4, 6 or 12 **AI**
 the order cannot be 1, 2, 3 or 6, since $a^6 \neq e$ **RI**
 the order cannot be 4, since $a^4 \neq e$ **RI**
 so the order of a must be 12
 therefore, a is a generator of G , which must therefore be cyclic **RI**
 another generator is eg a^{-1}, a^5, \dots **AI**

- (ii) $H = \{e, a^4, a^8\}$ **(A1)**

	e	a^4	a^8
e	e	a^4	a^8
a^4	a^4	a^8	e
a^8	a^8	e	a^4

M1A1
[9 marks]

- (b) no **AI**
 eg the group of symmetries of a triangle S_3 is not cyclic but all its (proper) subgroups are cyclic
 eg the Klein four-group is not cyclic but all its (proper) subgroups are cyclic **RI**

[2 marks]

Total [11 marks]

3. (a) (i)
$$\frac{1}{a^2+b^2} \begin{pmatrix} a & -b & 0 \\ b & a & 0 \\ 0 & 0 & a^2+b^2 \end{pmatrix} \begin{pmatrix} a & b & 0 \\ -b & a & 0 \\ 0 & 0 & 1 \end{pmatrix} = \mathbf{MI}$$

$$= \frac{1}{a^2+b^2} \begin{pmatrix} a^2+b^2 & 0 & 0 \\ 0 & a^2+b^2 & 0 \\ 0 & 0 & a^2+b^2 \end{pmatrix}$$
 which is the 3×3 identity \mathbf{AI}

Note: Award **AI** if it is clear the candidate recognizes this is the identity.

therefore
$$\begin{pmatrix} a & b & 0 \\ -b & a & 0 \\ 0 & 0 & 1 \end{pmatrix}^{-1} = \frac{1}{a^2+b^2} \begin{pmatrix} a & -b & 0 \\ b & a & 0 \\ 0 & 0 & a^2+b^2 \end{pmatrix} \mathbf{AG}$$

(ii)
$$A = \left\{ \begin{pmatrix} a & b & 0 \\ -b & a & 0 \\ 0 & 0 & 1 \end{pmatrix} \mid a, b \in \mathbb{R}, a^2+b^2 \neq 0 \right\}$$

attempt to show closure \mathbf{MI}

let $\begin{pmatrix} a & b & 0 \\ -b & a & 0 \\ 0 & 0 & 1 \end{pmatrix}$ and $\begin{pmatrix} c & d & 0 \\ -d & c & 0 \\ 0 & 0 & 1 \end{pmatrix}$ be two elements of A , then

$$\begin{pmatrix} a & b & 0 \\ -b & a & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} c & d & 0 \\ -d & c & 0 \\ 0 & 0 & 1 \end{pmatrix} \mathbf{(MI)}$$

$$= \begin{pmatrix} ac-bd & ad+bc & 0 \\ -bc-ad & -bd+ac & 0 \\ 0 & 0 & 1 \end{pmatrix} \mathbf{AI}$$

$$= \begin{pmatrix} f & g & 0 \\ -g & f & 0 \\ 0 & 0 & 1 \end{pmatrix},$$
 where $f = ac - bd$ and $g = ad + bc$ and

$$f^2 + g^2 (= (a^2+b^2)(c^2+d^2)) \neq 0 \mathbf{RI}$$

therefore A is closed under matrix multiplication

associativity follows from the fact that matrix multiplication is an associative binary operation on the collection of all 3×3 matrices \mathbf{AI}

continued ...

Question 3 continued

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \in A \quad \mathbf{AI}$$

for any matrix in A , $\det \begin{pmatrix} a & b & 0 \\ -b & a & 0 \\ 0 & 0 & 1 \end{pmatrix} = a^2 + b^2 \neq 0$ \mathbf{MI}

therefore, by part (i), it always has inverse

$$\begin{pmatrix} a & b & 0 \\ -b & a & 0 \\ 0 & 0 & 1 \end{pmatrix}^{-1} = \frac{1}{a^2 + b^2} \begin{pmatrix} a & -b & 0 \\ b & a & 0 \\ 0 & 0 & a^2 + b^2 \end{pmatrix} \in A \quad \mathbf{AI}$$

hence (A, \times) is a group \mathbf{AG}

[10 marks]

(b) consider the mapping (bijection) $f : A \rightarrow B$ defined by

$$f \left(\begin{pmatrix} a & b & 0 \\ -b & a & 0 \\ 0 & 0 & 1 \end{pmatrix} \right) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & a & -b \\ 0 & b & a \end{pmatrix} \quad \mathbf{(MI)}$$

$$f \left(\begin{pmatrix} a & b & 0 \\ -b & a & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} c & d & 0 \\ -d & c & 0 \\ 0 & 0 & 1 \end{pmatrix} \right) = f \left(\begin{pmatrix} ac - bd & ad + bc & 0 \\ -(ad + bc) & ac - bd & 0 \\ 0 & 0 & 1 \end{pmatrix} \right) \quad \mathbf{MI}$$

$$= \begin{pmatrix} 1 & 0 & 0 \\ 0 & ac - bd & -(ad + bc) \\ 0 & ad + bc & ac - bd \end{pmatrix} \quad \mathbf{AI}$$

$$f \left(\begin{pmatrix} a & b & 0 \\ -b & a & 0 \\ 0 & 0 & 1 \end{pmatrix} \right) f \left(\begin{pmatrix} c & d & 0 \\ -d & c & 0 \\ 0 & 0 & 1 \end{pmatrix} \right) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & a & -b \\ 0 & b & a \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & c & -d \\ 0 & d & c \end{pmatrix}$$

$$= \begin{pmatrix} 1 & 0 & 0 \\ 0 & ac - bd & -(ad + bc) \\ 0 & ad + bc & ac - bd \end{pmatrix} \quad \mathbf{AI}$$

therefore $f \left(\begin{pmatrix} a & b & 0 \\ -b & a & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} c & d & 0 \\ -d & c & 0 \\ 0 & 0 & 1 \end{pmatrix} \right) = f \left(\begin{pmatrix} a & b & 0 \\ -b & a & 0 \\ 0 & 0 & 1 \end{pmatrix} \right) f \left(\begin{pmatrix} c & d & 0 \\ -d & c & 0 \\ 0 & 0 & 1 \end{pmatrix} \right)$ \mathbf{RI}

[5 marks]

Total [15 marks]

4. (a) R is reflexive as $x^{-1} * x = e \in H \Rightarrow xRx$ for any $x \in G$ **AI**
 if xRy then $y^{-1} * x = h \in H$
 but $h \in H \Rightarrow h^{-1} \in H$, ie, $\underbrace{(y^{-1} * x)^{-1}}_{x^{-1} * y} \in H$ **MI**
- therefore yRx
 R is symmetric **AI**
 if xRy then $y^{-1} * x = h \in H$ and if yRz then $z^{-1} * y = k \in H$ **MI**
 $k * h \in H$, ie, $\underbrace{(z^{-1} * y) * (y^{-1} * x)}_{z^{-1} * x} \in H$ **AI**
- therefore xRz
 R is transitive **AI**
 so R is an equivalence relation on G **AG**
- [6 marks]**
- (b) $xRe \Leftrightarrow e^{-1} * x \in H$ **MI**
 $x \in H$ **AI**
 $[e] = H$ **AI** **N0**
- [3 marks]**

Total [9 marks]

5. (a) $(A \setminus B) \cup (B \setminus A) = (A \cap B') \cup (B \cap A')$ **(MI)**
 $= ((A \cap B') \cup B) \cap ((A \cap B') \cup A')$ **MI**
- $= \left((A \cup B) \cap \underbrace{(B' \cup B)}_U \right) \cap \left(\underbrace{(A \cup A')}_U \cap (B' \cup A') \right)$ **AI**
- $= (A \cup B) \cap (B' \cup A') = (A \cup B) \cap (A \cap B)'$ **AI**
 $= (A \cup B) \setminus (A \cap B)$ **AG**
- [4 marks]**

continued ...

Question 5 continued

(b) (i) false ***AI***
 counterexample
 eg $D \setminus C = \{2\} \notin S$ ***RI***

(ii) true ***AI***
 as $A \cap D = A$, $B \cap D = B$, $C \cap D = C$ and $D \cap D = D$,
 D is the identity ***RI***
 A (or B or C) has no inverse as $A \cap X = D$ is impossible ***RI***

(iii) false ***AI***
 when $Y = D$ the equation has more than one solution (four solutions) ***RI***

[7 marks]

Total [11 marks]