



MARKSCHEME

November 2006

MATHEMATICS

Standard Level

Paper 2

*This markscheme is **confidential** and for the exclusive use of examiners in this examination session.*

*It is the property of the International Baccalaureate and must **not** be reproduced or distributed to any other person without the authorization of IBCA.*

Instructions to Examiners

Abbreviations

- M** Marks awarded for attempting to use a correct **Method**; working must be seen.
- (M)** Marks awarded for **Method**; may be implied by **correct** subsequent working.
- A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- (A)** Marks awarded for an **Answer** or for **Accuracy**; may be implied by **correct** subsequent working.
- R** Marks awarded for clear **Reasoning**.
- N** Marks awarded for **correct** answers if **no** working shown.
- AG** Answer given in the question and so no marks are awarded.

Using the markscheme

1 General

Write the marks in red on candidates' scripts, in the right hand margin.

- Show the **breakdown** of individual marks awarded using the abbreviations **MI**, **AI**, etc.
- Write down the total for each **question** (at the end of the question) and **circle** it.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is not possible to award **M0** followed by **AI**, as **A** mark(s) depend on the preceding **M** mark(s), if any.
- Where **M** and **A** marks are noted on the same line, e.g. **MIAI**, this usually means **MI** for an **attempt** to use an appropriate method (e.g. substitution into a formula) and **AI** for using the **correct** values.
- Where the markscheme specifies (**M2**), **N3**, etc., do **not** split the marks.
- Once a correct answer to a question or part-question is seen, ignore further working.

3 N marks

*Award N marks for **correct** answers where there is **no** working.*

- Do **not** award a mixture of **N** and other marks.
- There may be fewer **N** marks available than the total of **M**, **A** and **R** marks; this is deliberate as it penalizes candidates for not following the instruction to show their working.
- For consistency within the markscheme, **N** marks are noted for every part, even when these match the mark breakdown.
- If a candidate has incorrect working, which somehow results in a correct answer, do **not** award the **N** marks for this correct answer. However, if the candidate has indicated (usually by crossing out) that the working is to be ignored, award the **N** marks for the correct answer.

4 Implied marks

*Implied marks appear in **brackets** e.g. (M1), and can only be awarded if **correct** work is seen or if implied in subsequent working.*

- Normally the correct work is seen or implied in the next line.
- Marks **without** brackets can only be awarded for work that is **seen**.

5 Follow through marks

*Follow through (FT) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s). To award FT marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part.*

- If the question becomes much simpler because of an error then use discretion to award fewer FT marks.
- If the error leads to an inappropriate value (e.g. probability greater than 1, use of $r > 1$ for the sum of an infinite GP, $\sin \theta = 1.5$), do not award the mark(s) for the final answer(s).
- Within a question part, once an error is made, no further A marks can be awarded, but M marks may be awarded if appropriate.
- Exceptions to this rule will be explicitly noted on the markscheme.

6 Mis-read

If a candidate incorrectly copies information from the question, this is a mis-read (MR). Apply a MR penalty of 1 mark to that question. Award the marks as usual and then write $-1(\text{MR})$ next to the total. Subtract 1 mark from the total for the question. A candidate should be penalized only once for a particular mis-read.

- If the question becomes much simpler because of the MR, then use discretion to award fewer marks.
- If the MR leads to an inappropriate value (e.g. $\sin \theta = 1.5$), do not award the mark(s) for the final answer(s).

7 Discretionary marks (d)

*An examiner uses discretion to award a mark on the rare occasions when the markscheme does not cover the work seen. The mark should be labelled (d) and a brief **note** written next to the mark explaining this decision.*

8 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If in doubt, contact your team leader for advice.

- Alternative methods for complete questions are indicated by **METHOD 1, METHOD 2, etc.**
- Alternative solutions for part-questions are indicated by **EITHER . . . OR.**
- Where possible, alignment will also be used to assist examiners in identifying where these alternatives start and finish.

9 Alternative forms

Unless the question specifies otherwise, **accept** equivalent forms.

- As this is an international examination, accept all alternative forms of **notation**.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer. This sometimes includes decimal/fraction equivalents.
- In the markscheme, **simplified** answers, (which candidates often do not write in examinations), will generally appear in brackets. Marks should be awarded for either the form preceding the bracket or the form in brackets (if it is seen).

Example: for differentiating $f(x) = 2 \sin(5x - 3)$, the markscheme gives:

$$f'(x) = (2 \cos(5x - 3)) 5 \quad (= 10 \cos(5x - 3)) \quad \text{AI}$$

Award **AI** for $(2 \cos(5x - 3)) 5$, even if $10 \cos(5x - 3)$ is not seen.

10 Accuracy of Answers

If the level of accuracy is specified in the question, a mark will be allocated for giving the answer to the required accuracy.

- **Rounding errors:** only applies to final answers not to intermediate steps.
- **Level of accuracy:** when this is not specified in the question the general rule applies: *unless otherwise stated in the question all numerical answers must be given exactly or correct to three significant figures.*

Candidates should be penalized **once only IN THE PAPER** for an accuracy error (**AP**). Award the marks as usual then write (**AP**) against the answer. On the **front** cover write $-1(\text{AP})$. Deduct 1 mark from the total for the paper, not the question.

- If a final correct answer is incorrectly rounded, apply the **AP**.
- If the level of accuracy is not specified in the question, apply the **AP** for correct answers not given to three significant figures.
- Intermediate values are sometimes written as 3.24(741). This indicates that using 3.24 (or 3.25) is acceptable, but the more accurate value is 3.24741. The digits in brackets are not required for the marks. If candidates work with fewer than three significant figures, this could lead to an **AP**.

If there is no working shown, and answers are given to the correct two significant figures, apply the **AP**. However, do **not** accept answers to one significant figure without working.

11 Crossed out work

If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work.

12 Examples

Exemplar material is available in the examiner training on Webct. Please refer to this material before you start marking, and when you have any queries. Please also feel free to contact your Team Leader if you need further advice.

QUESTION 1

Part A

- (a) (i) 50 (accept 49, “fewer than 50”) AI NI
- (ii) Cumulative frequency (7) = 90 (AI)
 $90 - 50$ (MI)
 $= 40$ AI N2
- (iii) 75th or 75.5th person AI
 median = 6.25 (min), 6 min 15 secs AI NI
- [6 marks]**

- (b) Evidence of finding 40 % (60 %) of 150 MI
 Number spending less than k minutes is $(150 - 60) = 90$ (AI)
 $k = 7$ AI N2
- [3 marks]**

- (c) (i)

t (minutes)	$0 \leq t < 2$	$2 \leq t < 4$	$4 \leq t < 6$	$6 \leq t < 8$	$8 \leq t < 10$	$10 \leq t < 12$
Frequency	10	23	37	38	27	15

AIAIAI N3

- (ii) Evidence of using **all** correct mid-interval values (1, 3, 5, 7, 9, 11) AI
- $$\text{mean} = \left(\frac{1 \times 10 + 3 \times 23 + 5 \times 37 + 7 \times 38 + 9 \times 27 + 11 \times 15}{150} \right)$$
- $= 6.25$ (min), 6 min 15 secs AI NI
- [5 marks]**

Sub-total [14 marks]

continued ...

Question 1 continued

Part B


- (a) (i) Attempt to set up sample space,
Any **correct** representation with 16 pairs

(MI)

A2

N3

e.g. 1,1 2,1 3,1 4,1
1,2 2,2 3,2 4,2
1,3 2,3 3,3 4,3
1,4 2,4 3,4 4,4



- (ii) Probability of two 4s is $\frac{1}{16}$ (= 0.0625)

A1

N1

[4 marks]

(b)

x	0	1	2
$P(X = x)$	$\frac{9}{16}$	$\frac{6}{16}$	$\frac{1}{16}$

A1A1A1

N3

[3 marks]

- (c) Evidence of selecting appropriate formula for $E(X)$

(MI)

e.g. $E(X) = \sum_0^2 xP(X = x)$, $E(X) = np$

Correct substitution

A1

e.g. $E(X) = 0 \times \frac{9}{16} + 1 \times \frac{6}{16} + 2 \times \frac{1}{16}$, $E(X) = 2 \times \frac{1}{4}$

$$E(X) = \frac{8}{16} \left(= \frac{1}{2} \right)$$

A1

N2

[3 marks]

Sub-total [10 marks]

Total [24 marks]

QUESTION 2

(a) $q = 0$ *AI* *NI*
[1 mark]

(b) Attempting to substitute (3, 18) *(MI)*
 $m^3 + n^3 + p^3 = 18$ *AI*
 $27m + 9n + 3p = 18$ *AG* *N0*
[2 marks]

(c) $m + n + p = 0$ *AI* *NI*
 $-m + n - p = -10$ *AI* *NI*
[2 marks]

(d) (i) Evidence of attempting to set up a matrix equation *(MI)*
 Correct **matrix** equation representing the given equations *A2* *N3*

$$e.g. \begin{pmatrix} 27 & 9 & 3 \\ 1 & 1 & 1 \\ -1 & 1 & -1 \end{pmatrix} \begin{pmatrix} m \\ n \\ p \end{pmatrix} = \begin{pmatrix} 18 \\ 0 \\ -10 \end{pmatrix}$$

(ii) $\begin{pmatrix} 2 \\ -5 \\ 3 \end{pmatrix}$ *AIAIAI* *N3*
[6 marks]

(e) Factorising *(MI)*
 $e.g. f(x) = x(2x^2 - 5x + 3)$, $f(x) = (x^2 - x)(rx - s)$
 $r = 2$ $s = 3$ (accept $f(x) = x(x - 1)(2x - 3)$) *AIAI* *N3*
[3 marks]

Total [14 marks]

QUESTION 3


Note: Throughout this question, the first and last terms are interchangeable.

- (a) For recognizing the arithmetic sequence *(MI)*
 $u_1 = 1, n = 20, u_{20} = 20$ ($u_1 = 1, n = 20, d = 1$) *(AI)*
 Evidence of using sum of an AP *MI*
 $S_{20} = \frac{(1+20)20}{2}$ (or $S = \frac{20}{2}(2 \times 1 + 19 \times 1)$) *AI*
 $S_{20} = 210$ *AG* *N0*
[4 marks]
- (b) Let there be n cans in bottom row *(MI)*
 Evidence of using $S_n = 3240$
 e.g. $\frac{(1+n)n}{2} = 3240, \frac{n}{2}(2 + (n-1)) = 3240, \frac{n}{2}(2n + (n-1)(-1)) = 3240$
 $n^2 + n - 6480 = 0$ *AI*
 $n = 80$ or $n = -81$ *(AI)*
 $n = 80$ *AI* *N2*
[4 marks]
- (c) (i) Evidence of using $S = \frac{(1+n)n}{2}$ *(MI)*
 $2S = n^2 + n$ *AI*
 $n^2 + n - 2S = 0$ *AG* *N0*
- (ii) **METHOD 1**
 Substituting $S = 2100$
 e.g. $n^2 + n - 4200 = 0, 2100 = \frac{(1+n)n}{2}$ *AI*
EITHER
 $n = 64.3, n = -65.3$ *AI*
 Any valid reason which includes reference to integer being needed, *RI*
 and pointing out that integer not possible here. *RI* *NI*
 e.g. n must be a (positive) integer, this equation does not have integer solutions.
- OR**
 Discriminant = 16 801 *AI*
 Valid reason which includes reference to integer being needed, *RI*
 and pointing out that integer not possible here. *RI* *NI*
 e.g. this discriminant is not a perfect square, therefore no integer solution as needed.
- METHOD 2**
 Trial and error
 $S_{64} = 2080, S_{65} = 2145$ *AIAI*
 Any valid reason which includes reference to integer being needed, *RI*
 and pointing out that integer not possible here. *RI* *NI*
[6 marks]

Total [14 marks]

QUESTION 4

(a)



AIAIAI

N3

Note: Award *AI* for the shape of the curve,
AI for correct domain,
AI for labelling **both** points P and Q in approximately correct positions.

[3 marks]

(b) (i) Correctly finding derivative of $2x+1$ i.e. 2 (AI)
 Correctly finding derivative of e^{-x} i.e. $-e^{-x}$ (AI)
 Evidence of using the product rule (MI)
 $f'(x) = 2e^{-x} + (2x+1)(-e^{-x})$ AI
 $= (1-2x)e^{-x}$ AG N0

(ii) At Q, $f'(x) = 0$ (MI)
 $x = 0.5, y = 2e^{-0.5}$ AIAI
 Q is $(0.5, 2e^{-0.5})$ N3

[7 marks]

(c) $1 \leq k < 2e^{-0.5}$ A2 N2
 [2 marks]

(d) Using $f''(x) = 0$ at the point of inflexion MI
 $e^{-x}(-3+2x) = 0$
 This equation has only one root. RI
 So f has only one point of inflexion. AG N0
 [2 marks]

continued ...

Question 4 continued

(e) At R, $y = 7e^{-3}$ ($= 0.34850 \dots$) **(A1)**

Gradient of (PR) is $\frac{7e^{-3}-1}{3}$ ($= -0.2172$) **(A1)**

Equation of (PR) is $g(x) = \left(\frac{7e^{-3}-1}{3}\right)x + 1$ ($= -0.2172x + 1$) **A1**

Evidence of appropriate method, involving subtraction of integrals or areas **M2**

Correct limits/endpoints **A1**

e.g. $\int_0^3 (f(x) - g(x)) dx$, area under curve – area under PR

$$\text{Shaded area is } \int_0^3 \left((2x+1)e^{-x} - \left(\frac{7e^{-3}-1}{3}x + 1 \right) \right) dx$$

$$= 0.529$$

A1 **N4**
[7 marks]

Total [21 marks]

QUESTION 5

- (a) (i) $OP = PQ$ (= 3 cm) **RI**
 So $\triangle OPQ$ is isosceles **AG** **N0**
- (ii) Using cos rule correctly *e.g.* $\cos \hat{O}PQ = \frac{3^2 + 3^2 - 4^2}{2 \times 3 \times 3}$ **(MI)**
 $\cos \hat{O}PQ = \frac{9 + 9 - 16}{18} \left(= \frac{2}{18} \right)$ **AI**
 $\cos \hat{O}PQ = \frac{1}{9}$ **AG** **N0**
- (iii) Evidence of using $\sin^2 A + \cos^2 A = 1$ **MI**
 $\sin \hat{O}PQ = \sqrt{1 - \frac{1}{81}} \left(= \sqrt{\frac{80}{81}} \right)$ **AI**
 $\sin \hat{O}PQ = \frac{\sqrt{80}}{9}$ **AG** **N0**
- (iv) Evidence of using area triangle $OPQ = \frac{1}{2} \times OP \times PQ \sin P$ **MI**
e.g. $\frac{1}{2} \times 3 \times 3 \times \frac{\sqrt{80}}{9}, \frac{9}{2} \times 0.9938...$
 Area triangle $OPQ = \frac{\sqrt{80}}{2} \left(= \sqrt{20} \right) (= 4.47)$ **AI** **N1**
[7 marks]
- (b) (i) $\hat{O}PQ = 1.4594...$
 $\hat{O}PQ = 1.46$ **AI** **N1**
- (ii) Evidence of using formula for area of a sector **(MI)**
e.g. Area sector $OPQ = \frac{1}{2} \times 3^2 \times 1.4594...$
 $= 6.57$ **AI** **N2**
[3 marks]
- (c) $\hat{Q}OP = \frac{\pi - 1.4594...}{2} (= 0.841)$ **(AI)**
 Area sector $QOS = \frac{1}{2} \times 4^2 \times 0.841$ **AI**
 $= 6.73$ **AI** **N2**
[3 marks]

continued ...

Question 5 continued

(d) Area of small semicircle is 4.5π (=14.137...)

AI

Evidence of correct approach

MI

e.g. Area = area of semicircle – area sector OPQ – area sector QOS + area triangle POQ

Correct expression

AI

e.g. $4.5\pi - 6.5675... - 6.7285... + 4.472...$, $4.5\pi - (6.7285... + 2.095...)$,
 $4.5\pi - (6.5675... + 2.256...)$

Area of the shaded region = 5.31

AI

NI

[4 marks]

Total [17 marks]
