



MARKSCHEME

May 2010

MATHEMATICS

Higher Level

Paper 1

Samples to team leaders	June 7 2010
Everything (marks, scripts etc) to IB Cardiff	June 14 2010

*This markscheme is **confidential** and for the exclusive use of examiners in this examination session.*

*It is the property of the International Baccalaureate and must **not** be reproduced or distributed to any other person without the authorization of IB Cardiff.*

Instructions to Examiners

Abbreviations

- M** Marks awarded for attempting to use a correct **Method**; working must be seen.
- (M)** Marks awarded for **Method**; may be implied by **correct** subsequent working.
- A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- (A)** Marks awarded for an **Answer** or for **Accuracy**; may be implied by **correct** subsequent working.
- R** Marks awarded for clear **Reasoning**.
- N** Marks awarded for **correct** answers if **no** working shown.
- AG** Answer given in the question and so no marks are awarded.

Using the markscheme

1 General

Write the marks in red on candidates' scripts, in the right hand margin.

- Show the **breakdown** of individual marks awarded using the abbreviations **MI**, **AI**, etc.
- Write down the total for each **question** (at the end of the question) and **circle** it.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is not possible to award **M0** followed by **AI**, as **A** mark(s) depend on the preceding **M** mark(s), if any.
- Where **M** and **A** marks are noted on the same line, e.g. **MIAI**, this usually means **MI** for an **attempt** to use an appropriate method (e.g. substitution into a formula) and **AI** for using the **correct** values.
- Where the markscheme specifies (**M2**), **N3**, etc., do **not** split the marks.
- Once a correct answer to a question or part-question is seen, ignore further working.

3 N marks

*Award N marks for **correct** answers where there is **no** working.*

- Do **not** award a mixture of **N** and other marks.
- There may be fewer **N** marks available than the total of **M**, **A** and **R** marks; this is deliberate as it penalizes candidates for not following the instruction to show their working.

4 Implied marks

*Implied marks appear in **brackets e.g. (M1)**, and can only be awarded if **correct** work is seen or if implied in subsequent working.*

- Normally the correct work is seen or implied in the next line.
- Marks **without** brackets can only be awarded for work that is **seen**.

5 Follow through marks

*Follow through (**FT**) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s). To award **FT** marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part.*

- If the question becomes much simpler because of an error then use discretion to award fewer **FT** marks.
- If the error leads to an inappropriate value (e.g. $\sin \theta = 1.5$), do not award the mark(s) for the final answer(s).
- Within a question part, once an error is made, no further **dependent A** marks can be awarded, but **M** marks may be awarded if appropriate.
- Exceptions to this rule will be explicitly noted on the markscheme.

6 Mis-read

*If a candidate incorrectly copies information from the question, this is a mis-read (**MR**). Apply a **MR** penalty of 1 mark to that question. Award the marks as usual and then write $-1(\mathbf{MR})$ next to the total. Subtract 1 mark from the total for the question. A candidate should be penalized only once for a particular mis-read.*

- If the question becomes much simpler because of the **MR**, then use discretion to award fewer marks.
- If the **MR** leads to an inappropriate value (e.g. $\sin \theta = 1.5$), do not award the mark(s) for the final answer(s).

7 Discretionary marks (*d*)

*An examiner uses discretion to award a mark on the rare occasions when the markscheme does not cover the work seen. The mark should be labelled (**d**) and a brief **note** written next to the mark explaining this decision.*

8 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If in doubt, contact your team leader for advice.

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, etc.
- Alternative solutions for part-questions are indicated by **EITHER . . . OR**.
- Where possible, alignment will also be used to assist examiners in identifying where these alternatives start and finish.

9 Alternative forms

Unless the question specifies otherwise, **accept** equivalent forms.

- As this is an international examination, accept all alternative forms of **notation**.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, **simplified** answers, (which candidates often do not write in examinations), will generally appear in brackets. Marks should be awarded for either the form preceding the bracket or the form in brackets (if it is seen).

Example: for differentiating $f(x) = 2\sin(5x - 3)$, the markscheme gives:

$$f'(x) = 2\cos(5x - 3) \cdot 5 = 10\cos(5x - 3) \quad \mathbf{AI}$$

Award **AI** for $2\cos(5x - 3) \cdot 5$, even if $10\cos(5x - 3)$ is not seen.

10 Accuracy of Answers

If the level of accuracy is specified in the question, a mark will be allocated for giving the answer to the required accuracy.

- **Rounding errors:** only applies to final answers not to intermediate steps.
- **Level of accuracy:** when this is not specified in the question the general rule applies: *unless otherwise stated in the question all numerical answers must be given exactly or correct to three significant figures.*

Candidates should be penalized **once only IN THE PAPER** for an accuracy error (**AP**). Award the marks as usual then write (**AP**) against the answer. On the **front** cover write $-1(\mathbf{AP})$. Deduct 1 mark from the total for the paper, not the question.

- If a final correct answer is incorrectly rounded, apply the **AP**.
- If the level of accuracy is not specified in the question, apply the **AP** for correct answers not given to three significant figures.

If there is no working shown, and answers are given to the correct two significant figures, apply the **AP**. However, do **not** accept answers to one significant figure without working.

11 Crossed out work

If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work.

SECTION A

1. (a) the total area under the graph of the pdf is unity (AI)

$$\text{area} = c \int_0^1 x - x^2 \, dx$$

$$= c \left[\frac{1}{2}x^2 - \frac{1}{3}x^3 \right]_0^1 \quad \text{AI}$$

$$= \frac{c}{6}$$

$$\Rightarrow c = 6 \quad \text{AI}$$

- (b) $E(X) = 6 \int_0^1 x^2 - x^3 \, dx$ (MI)

$$= 6 \left(\frac{1}{3} - \frac{1}{4} \right) = \frac{1}{2} \quad \text{AI}$$

Note: Allow an answer obtained by a symmetry argument.

[5 marks]

2. (a) attempt at completing the square (MI)

$$3x^2 - 6x + 5 = 3(x^2 - 2x) + 5 = 3(x - 1)^2 - 1 + 5 \quad \text{(AI)}$$

$$= 3(x - 1)^2 + 2 \quad \text{AI}$$

$$(a = 3, b = -1, c = 2)$$

- (b) definition of suitable basic transformations:
 $T_1 =$ stretch in y direction scale factor 3 AI

$$T_2 = \text{translation} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad \text{AI}$$

$$T_3 = \text{translation} \begin{pmatrix} 0 \\ 2 \end{pmatrix} \quad \text{AI}$$

[6 marks]

3. (a) $2y + 8x = 4$ *MI*
 $-3x + 2y = -7$ *AI*
 $2x + 6 - 2x = 6$

Note: Award *MI* for attempt at components, *AI* for two correct equations.
 No penalty for not checking the third equation.

solving : $x = 1, y = -2$ *AI*

(b) $|a + 2b| = \left| \begin{pmatrix} -4 \\ -3 \\ 2 \end{pmatrix} + 2 \begin{pmatrix} 4 \\ -2 \\ 2 \end{pmatrix} \right|$
 $= \left| \begin{pmatrix} 4 \\ -7 \\ 6 \end{pmatrix} \right|$

$\Rightarrow |a + 2b| = \sqrt{4^2 + (-7)^2 + 6^2}$ *(MI)*

$= \sqrt{101}$ *AI*

[5 marks]

4. recognition of $X \sim B\left(6, \frac{4}{7}\right)$ *(MI)*

$P(X = 3) = \binom{6}{3} \left(\frac{4}{7}\right)^3 \left(\frac{3}{7}\right)^3 = 20 \times \frac{4^3 \times 3^3}{7^6}$ *AI*

$P(X = 2) = \binom{6}{2} \left(\frac{4}{7}\right)^2 \left(\frac{3}{7}\right)^4 = 15 \times \frac{4^2 \times 3^4}{7^6}$ *AI*

$\frac{P(X = 3)}{P(X = 2)} = \frac{80}{45} = \frac{16}{9}$ *AI*

[4 marks]

5. (a) $BA = \left(\begin{pmatrix} 1 & 3 \\ 2 & -2 \end{pmatrix} \begin{pmatrix} 3 & -2 \\ 5 & -4 \end{pmatrix} \right) = \begin{pmatrix} 18 & -14 \\ -4 & 4 \end{pmatrix}$ *A2*

Note: Award *AI* for one error, *A0* for two or more errors.

(b) $\det(BA) = (72 - 56) = 16$ *(MI)AI*

(c) **EITHER**

$A(A^{-1}B + 2A^{-1})A = BA + 2A$ *(MI)(AI)*

$$= \begin{pmatrix} 24 & -18 \\ 6 & -4 \end{pmatrix} \quad \text{AI}$$

OR

$A^{-1} = -\frac{1}{2} \begin{pmatrix} -4 & 2 \\ -5 & 3 \end{pmatrix}$ *(AI)*

an attempt to evaluate *(MI)*

$$A^{-1}B + 2A^{-1} = -\frac{1}{2} \begin{pmatrix} 0 & -16 \\ 1 & -21 \end{pmatrix} - \begin{pmatrix} -4 & 2 \\ -5 & 3 \end{pmatrix}$$

$$= \begin{pmatrix} 4 & 6 \\ 4.5 & 7.5 \end{pmatrix}$$

$A(A^{-1}B + 2A^{-1})A = \begin{pmatrix} 3 & -2 \\ 5 & -4 \end{pmatrix} \begin{pmatrix} 4 & 6 \\ 4.5 & 7.5 \end{pmatrix} \begin{pmatrix} 3 & -2 \\ 5 & -4 \end{pmatrix}$

$$= \begin{pmatrix} 3 & 3 \\ 2 & 0 \end{pmatrix} \begin{pmatrix} 3 & -2 \\ 5 & -4 \end{pmatrix} = \begin{pmatrix} 24 & -18 \\ 6 & -4 \end{pmatrix} \quad \text{AI}$$

[7 marks]

6. $\sin\left(x + \frac{\pi}{3}\right) = \sin x \cos\left(\frac{\pi}{3}\right) + \cos x \sin\left(\frac{\pi}{3}\right)$ (M1)

$$\sin x \cos\left(\frac{\pi}{3}\right) + \cos x \sin\left(\frac{\pi}{3}\right) = 2 \sin x \sin\left(\frac{\pi}{3}\right)$$

$$\frac{1}{2} \sin x + \frac{\sqrt{3}}{2} \cos x = 2 \times \frac{\sqrt{3}}{2} \sin x$$
 A1

dividing by $\cos x$ and rearranging M1

$$\tan x = \frac{\sqrt{3}}{2\sqrt{3}-1}$$
 A1

rationalizing the denominator M1

$$11 \tan x = 6 + \sqrt{3}$$
 A1

[6 marks]

7. (a) $\left(u = x^2 - 2x - 1.5; \frac{du}{dx} = 2x - 2\right)$

$$\frac{df}{dx} = \frac{df}{du} \frac{du}{dx} = e^u (2x - 2)$$
 (M1)

$$= 2(x-1)e^{x^2-2x-1.5}$$
 A1

(b) $\frac{dy}{dx} = \frac{(x-1) \times 2(x-1)e^{x^2-2x-1.5} - 1 \times e^{x^2-2x-1.5}}{(x-1)^2}$ M1A1

$$= \frac{2x^2 - 4x + 1}{(x-1)^2} e^{x^2-2x-1.5}$$
 (A1)

minimum occurs when $\frac{dy}{dx} = 0$ (M1)

$$x = 1 \pm \sqrt{\frac{1}{2}} \left(\text{accept } x = \frac{4 \pm \sqrt{8}}{4} \right)$$
 A1

$$a = 1 + \sqrt{\frac{1}{2}} \left(\text{accept } a = \frac{4 + \sqrt{8}}{4} \right)$$
 R1

[8 marks]

8. EITHER

differentiating implicitly:

$$1 \times e^{-y} - xe^{-y} \frac{dy}{dx} + e^y \frac{dy}{dx} = 1 \quad \text{MIAI}$$

at the point $(c, \ln c)$

$$\frac{1}{c} - c \times \frac{1}{c} \frac{dy}{dx} + c \frac{dy}{dx} = 1 \quad \text{MI}$$

$$\frac{dy}{dx} = \frac{1}{c} \quad (c \neq 1) \quad \text{(AI)}$$

OR

reasonable attempt to make expression explicit (MI)

$$xe^{-y} + e^y = 1 + x$$

$$x + e^{2y} = e^y (1 + x)$$

$$e^{2y} - e^y (1 + x) + x = 0$$

$$(e^y - 1)(e^y - x) = 0 \quad \text{(AI)}$$

$$e^y = 1, e^y = x$$

$$y = 0, y = \ln x \quad \text{AI}$$

Note: Do not penalize if $y = 0$ not stated.

$$\frac{dy}{dx} = \frac{1}{x}$$

$$\text{gradient of tangent} = \frac{1}{c} \quad \text{AI}$$

Note: If candidate starts with $y = \ln x$ with no justification, award (M0)(A0)AIAI.

THEN

the equation of the normal is

$$y - \ln c = -c(x - c) \quad \text{MI}$$

$$x = 0, y = c^2 + 1$$

$$c^2 + 1 - \ln c = c^2 \quad \text{(AI)}$$

$$\ln c = 1$$

$$c = e \quad \text{AI}$$

[7 marks]

9. EITHER

attempt at integration by substitution

*(M1)*using $u = t + 1$, $du = dt$, the integral becomes

$$\int_1^2 (u-1) \ln u \, du$$

AI

then using integration by parts

M1

$$\int_1^2 (u-1) \ln u \, du = \left[\left(\frac{u^2}{2} - u \right) \ln u \right]_1^2 - \int_1^2 \left(\frac{u^2}{2} - u \right) \times \frac{1}{u} \, du$$

AI

$$= - \left[\frac{u^2}{4} - u \right]_1^2$$

(AI)

$$= \frac{1}{4} \quad (\text{accept } 0.25)$$

*AI***OR**

attempt to integrate by parts

(M1)

correct choice of variables to integrate and differentiate

M1

$$\int_0^1 t \ln(t+1) \, dt = \left[\frac{t^2}{2} \ln(t+1) \right]_0^1 - \int_0^1 \frac{t^2}{2} \times \frac{1}{t+1} \, dt$$

AI

$$= \left[\frac{t^2}{2} \ln(t+1) \right]_0^1 - \frac{1}{2} \int_0^1 t - 1 + \frac{1}{t+1} \, dt$$

AI

$$= \left[\frac{t^2}{2} \ln(t+1) \right]_0^1 - \frac{1}{2} \left[\frac{t^2}{2} - t + \ln(t+1) \right]_0^1$$

(AI)

$$= \frac{1}{4} \quad (\text{accept } 0.25)$$

*AI**[6 marks]*

10. (a) **Note:** Interchange of variables may take place at any stage.

for the inverse, solve for x in

$$y = \frac{2x-3}{x-1}$$

$$y(x-1) = 2x-3$$

MI

$$yx - 2x = y - 3$$

$$x(y-2) = y-3$$

(AI)

$$x = \frac{y-3}{y-2}$$

$$\Rightarrow f^{-1}(x) = \frac{x-3}{x-2} \quad (x \neq 2)$$

AI

Note: Do not award final **AI** unless written in the form $f^{-1}(x) = \dots$

(b) $\pm f^{-1}(x) = 1 + f^{-1}(x)$ leads to

$$2 \frac{x-3}{x-2} = -1$$

(MI)AI

$$x = \frac{8}{3}$$

AI

[6 marks]

SECTION B

11. (a) (i) $1 \times 2 + 2 \times 3 + \dots + n(n+1) = \frac{1}{3}n(n+1)(n+2)$ **RI**
- (ii) LHS = 40; RHS = 40 **AI**
[2 marks]

- (b) the sequence of values are:
 5, 7, 11, 19, 35 ... or an example **AI**
 35 is not prime, so Bill's conjecture is false **RIAG**
[2 marks]

- (c) $P(n)$: $5 \times 7^n + 1$ is divisible by 6 **AI**
 $P(1)$: 36 is divisible by 6 $\Rightarrow P(1)$ true **AI**
 assume $P(k)$ is true ($5 \times 7^k + 1 = 6r$) **MI**

Note: Do **not** award **MI** for statement starting 'let $n = k$ '.
 Subsequent marks are independent of this **MI**.

- consider $5 \times 7^{k+1} + 1$ **MI**
 $= 7(6r - 1) + 1$ **(AI)**
 $= 6(7r - 1) \Rightarrow P(k+1)$ is true **AI**
 $P(1)$ true and $P(k)$ true $\Rightarrow P(k+1)$ true, so by **MI** $P(n)$ is true for all $n \in \mathbb{Z}^+$ **RI**

Note: Only award **RI** if there is consideration of $P(1)$, $P(k)$ and $P(k+1)$ in the final statement.

Only award **RI** if at least one of the two preceding **A** marks has been awarded.

[6 marks]

Total [10 marks]

12. (a) (i) use of $\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}| |\mathbf{b}| \cos \theta$ (MI)
 $\mathbf{a} \cdot \mathbf{b} = -1$ (AI)
 $|\mathbf{a}| = 7, |\mathbf{b}| = 5$ (AI)
 $\cos \theta = -\frac{1}{35}$ AI

- (ii) the required cross product is

$$\begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 6 & 3 & 2 \\ 0 & -3 & 4 \end{vmatrix} = 18\mathbf{i} - 24\mathbf{j} - 18\mathbf{k}$$
 MIAI

- (iii) using $\mathbf{r} \cdot \mathbf{n} = \mathbf{p} \cdot \mathbf{n}$ the equation of the plane is (MI)
 $18x - 24y - 18z = 12$ ($3x - 4y - 3z = 2$) AI

- (iv) recognizing that $z = 0$ (MI)
 x -intercept = $\frac{2}{3}$, y -intercept = $-\frac{1}{2}$ (AI)
 $\text{area} = \left(\frac{2}{3}\right)\left(\frac{1}{2}\right)\left(\frac{1}{2}\right) = \frac{1}{6}$ AI

[11 marks]

- (b) (i) $\mathbf{p} \cdot \mathbf{p} = |\mathbf{p}| |\mathbf{p}| \cos 0$ MIAI
 $= |\mathbf{p}|^2$ AG

- (ii) consider the LHS, and use of result from part (i)
 $|\mathbf{p} + \mathbf{q}|^2 = (\mathbf{p} + \mathbf{q}) \cdot (\mathbf{p} + \mathbf{q})$ MI
 $= \mathbf{p} \cdot \mathbf{p} + \mathbf{p} \cdot \mathbf{q} + \mathbf{q} \cdot \mathbf{p} + \mathbf{q} \cdot \mathbf{q}$ (AI)
 $= \mathbf{p} \cdot \mathbf{p} + 2\mathbf{p} \cdot \mathbf{q} + \mathbf{q} \cdot \mathbf{q}$ AI
 $= |\mathbf{p}|^2 + 2\mathbf{p} \cdot \mathbf{q} + |\mathbf{q}|^2$ AG

continued ...

Question 12 continued

(iii) EITHER

use of $p \cdot q \leq |p||q|$

MI

so $0 \leq |p+q|^2 = |p|^2 + 2p \cdot q + |q|^2 \leq |p|^2 + 2|p||q| + |q|^2$

AI


take square root (of these positive quantities) to establish

AI

$|p+q| \leq |p| + |q|$

AG

OR



MIMI

Note: Award *MI* for correct diagram and *MI* for correct labelling of vectors including arrows.

since the sum of any two sides of a triangle is greater than the third side,

$|p| + |q| > |p+q|$

AI

when p and q are collinear $|p| + |q| = |p+q|$

$\Rightarrow |p+q| \leq |p| + |q|$

AG

[8 marks]


Total [19 marks]

13. (a) (i) $\omega^3 = \left(\cos\left(\frac{2\pi}{3}\right) + i \sin\left(\frac{2\pi}{3}\right) \right)^3$
 $= \cos\left(3 \times \frac{2\pi}{3}\right) + i \sin\left(3 \times \frac{2\pi}{3}\right)$ *(M1)*
 $= \cos 2\pi + i \sin 2\pi$ *AI*
 $= 1$ *AG*

(ii) $1 + \omega + \omega^2 = 1 + \cos\left(\frac{2\pi}{3}\right) + i \sin\left(\frac{2\pi}{3}\right) + \cos\left(\frac{4\pi}{3}\right) + i \sin\left(\frac{4\pi}{3}\right)$ *M1A1*
 $= 1 + \frac{1}{2} + i \frac{\sqrt{3}}{2} - \frac{1}{2} - i \frac{\sqrt{3}}{2}$ *AI*
 $= 0$ *AG*

[5 marks]

(b) (i) $e^{i\theta} + e^{i\left(\theta + \frac{2\pi}{3}\right)} + e^{i\left(\theta + \frac{4\pi}{3}\right)}$
 $= e^{i\theta} + e^{i\theta} e^{i\left(\frac{2\pi}{3}\right)} + e^{i\theta} e^{i\left(\frac{4\pi}{3}\right)}$ *(M1)*
 $= \left(e^{i\theta} \left(1 + e^{i\left(\frac{2\pi}{3}\right)} + e^{i\left(\frac{4\pi}{3}\right)} \right) \right)$
 $= e^{i\theta} (1 + \omega + \omega^2)$ *AI*
 $= 0$ *AG*



A1A1

Note: Award *AI* for one point on the imaginary axis and another point marked with approximately correct modulus and argument. Award *AI* for third point marked to form an equilateral triangle centred on the origin.

[4 marks]

continued ...

Question 13 continued

- (c) (i) attempt at the expansion of at least two linear factors (MI)
 $(z-1) z^2 - z(\omega + \omega^2) + \omega^3$ or equivalent (AI)
 use of earlier result (MI)
 $F(z) = (z-1)(z^2 + z + 1) = z^3 - 1$ AI
- (ii) equation to solve is $z^3 = 8$ (MI)
 $z = 2, 2\omega, 2\omega^2$ A2

Note: Award **AI** for 2 correct solutions.

[7 marks]

Total [16 marks]

14. (a) the differential equation is separable and can be written as (MI)
 $\int -y^{-2} dy = \int \cos^2 x dx$ (or equivalent) AI
 $= \int \frac{1 + \cos 2x}{2} dx$ AI
 $\frac{1}{y} = \frac{1}{2}x + \frac{1}{4}\sin 2x (+C)$ AIAI
 when $x = 0, y = 1$ MI
 $C = 1$
 $y = \frac{1}{\frac{1}{2}x + \frac{1}{4}\sin 2x + 1}$ AI

[7 marks]

- (b) (i) recognizing use of $(1 + \tan x)^2$ (MI)
 $(1 + \tan x)^2 = 1 + 2 \tan x + \tan^2 x \geq 1 + \tan^2 x = \sec^2 x$ AI
 (since all terms are positive)
 $(1 + \tan x)^2 \geq \sec^2 x$
 $\sec^2 x = 1 + \tan^2 x \geq 1$ AI
 $\Rightarrow (1 + \tan x)^2 \geq \sec^2 x \geq 1$
 since all terms are positive, taking square root gives MI
 $1 \leq \sec x \leq 1 + \tan x$ AG

- (ii) $\int_0^{\frac{\pi}{4}} dx \leq \int_0^{\frac{\pi}{4}} \sec x dx \leq \int_0^{\frac{\pi}{4}} 1 + \tan x dx$ MI
 $x \Big|_0^{\frac{\pi}{4}} \leq \int_0^{\frac{\pi}{4}} \sec x dx \leq x - \ln \cos x \Big|_0^{\frac{\pi}{4}}$ MIAI
 $\frac{\pi}{4} \leq \int_0^{\frac{\pi}{4}} \sec x dx \leq \frac{\pi}{4} - \ln \frac{1}{\sqrt{2}}$ AI
 $\frac{\pi}{4} \leq \int_0^{\frac{\pi}{4}} \sec x dx \leq \frac{\pi}{4} + \frac{1}{2} \ln 2$ AG

[8 marks]

Total [15 marks]