



MARKSCHEME

May 2012

MATHEMATICS

Higher Level

Paper 1

*This markscheme is **confidential** and for the exclusive use of examiners in this examination session.*

*It is the property of the International Baccalaureate and must **not** be reproduced or distributed to any other person without the authorization of IB Cardiff.*

Instructions to Examiners

Abbreviations

- M** Marks awarded for attempting to use a correct **Method**; working must be seen.
- (M)** Marks awarded for **Method**; may be implied by **correct** subsequent working.
- A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- (A)** Marks awarded for an **Answer** or for **Accuracy**; may be implied by **correct** subsequent working.
- R** Marks awarded for clear **Reasoning**.
- N** Marks awarded for **correct** answers if **no** working shown.
- AG** Answer given in the question and so no marks are awarded.

Using the markscheme

1 General

Mark according to scoris instructions and the document “**Mathematics HL: Guidance for e-marking May 2012**”. It is essential that you read this document before you start marking. In particular, please note the following:

- Marks must be recorded using the annotation stamps. Please check that you are entering marks for the right question.
- If a part is **completely correct**, (and gains all the “must be seen” marks), use the ticks with numbers to stamp full marks.
- If a part is completely wrong, stamp **A0** by the final answer.
- If a part gains anything else, it **must** be recorded using **all** the annotations.
- All the marks will be added and recorded by scoris.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is not possible to award **M0** followed by **AI**, as **A** mark(s) depend on the preceding **M** mark(s), if any.
- Where **M** and **A** marks are noted on the same line, *e.g.* **MIAI**, this usually means **MI** for an **attempt** to use an appropriate method (*e.g.* substitution into a formula) and **AI** for using the **correct** values.
- Where the markscheme specifies (**M2**), **N3**, *etc.*, do **not** split the marks.
- Once a correct answer to a question or part-question is seen, ignore further working.

3 N marks

Award **N** marks for **correct** answers where there is **no** working.

- Do **not** award a mixture of **N** and other marks.
- There may be fewer **N** marks available than the total of **M**, **A** and **R** marks; this is deliberate as it penalizes candidates for not following the instruction to show their working.

4 Implied marks

*Implied marks appear in **brackets e.g. (M1)**, and can only be awarded if **correct** work is seen or if implied in subsequent working.*

- Normally the correct work is seen or implied in the next line.
- Marks **without** brackets can only be awarded for work that is **seen**.

5 Follow through marks

*Follow through (**FT**) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s). To award **FT** marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part.*

- If the question becomes much simpler because of an error then use discretion to award fewer **FT** marks.
- If the error leads to an inappropriate value (e.g. $\sin \theta = 1.5$), do not award the mark(s) for the final answer(s).
- Within a question part, once an error is made, no further **dependent A** marks can be awarded, but **M** marks may be awarded if appropriate.
- Exceptions to this rule will be explicitly noted on the markscheme.

6 Mis-read

*If a candidate incorrectly copies information from the question, this is a mis-read (**MR**). A candidate should be penalized only once for a particular mis-read. Use the **MR** stamp to indicate that this has been a misread. Then deduct the first of the marks to be awarded, even if this is an **M** mark, but award all others so that the candidate only loses one mark.*

- If the question becomes much simpler because of the **MR**, then use discretion to award fewer marks.
- If the **MR** leads to an inappropriate value (e.g. $\sin \theta = 1.5$), do not award the mark(s) for the final answer(s).

7 Discretionary marks (*d*)

*An examiner uses discretion to award a mark on the rare occasions when the markscheme does not cover the work seen. In such cases the annotation **DM** should be used and a brief **note** written next to the mark explaining this decision.*

8 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If in doubt, contact your team leader for advice.

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, etc.
- Alternative solutions for part-questions are indicated by **EITHER . . . OR**.
- Where possible, alignment will also be used to assist examiners in identifying where these alternatives start and finish.

9 Alternative forms

Unless the question specifies otherwise, **accept** equivalent forms.

- As this is an international examination, accept all alternative forms of **notation**.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, **simplified** answers, (which candidates often do not write in examinations), will generally appear in brackets. Marks should be awarded for either the form preceding the bracket or the form in brackets (if it is seen).

Example: for differentiating $f(x) = 2\sin(5x - 3)$, the markscheme gives:

$$f'(x) = (2\cos(5x - 3))5 \quad (=10\cos(5x - 3)) \quad \text{AI}$$

Award **AI** for $(2\cos(5x - 3))5$, even if $10\cos(5x - 3)$ is not seen.

10 Accuracy of Answers

Candidates should **NO LONGER** be penalized for an accuracy error (**AP**).

If the level of accuracy is specified in the question, a mark will be allocated for giving the answer to the required accuracy. When this is not specified in the question, all numerical answers should be given exactly or correct to three significant figures. Please check work carefully for **FT**.

11 Crossed out work

If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work.

12 Calculators

No calculator is allowed. The use of any calculator on paper 1 is malpractice, and will result in no grade awarded. If you see work that suggests a candidate has used any calculator, please follow the procedures for malpractice. Examples: finding an angle, given a trig ratio of 0.4235.

13 More than one solution

Where a candidate offers two or more different answers to the same question, an examiner should only mark the first response unless the candidate indicates otherwise.

SECTION A

1. $u_1 = \frac{1}{3}k, r = \frac{1}{3}$

(AI)(AI)

$$7 = \frac{\frac{1}{3}k}{1 - \frac{1}{3}}$$


MI

$k = 14$

AI

[4 marks]

2. (a)



MIAIAIAI

Note: Award *MI* for any of the three sections completely correct, *AI* for each correct segment of the graph.

[4 marks]

(b) (i) 0

AI

(ii) 2

AI

(iii) finding area of rectangle
-4

(MI)

AI

Note: Award *MIA0* for the answer 4.

[4 marks]

Total [8 marks]

3. $z_1 = 2a \operatorname{cis}\left(\frac{\pi}{3}\right), z_2 = \sqrt{2} \operatorname{cis}\left(-\frac{\pi}{4}\right)$

MIAIAI

EITHER

$$\left(\frac{z_1}{z_2}\right)^6 = \frac{2^6 a^6 \operatorname{cis}(0)}{\sqrt{2}^6 \operatorname{cis}\left(\frac{\pi}{2}\right)} \left(= 8a^6 \operatorname{cis}\left(-\frac{\pi}{2}\right)\right)$$

MIAIAI

OR

$$\begin{aligned} \left(\frac{z_1}{z_2}\right)^6 &= \left(\frac{2a}{\sqrt{2}} \operatorname{cis}\left(\frac{7\pi}{12}\right)\right)^6 \\ &= 8a^6 \operatorname{cis}\left(-\frac{\pi}{2}\right) \end{aligned}$$

MIAI

AI

THEN


$$= -8a^6 i$$

AI

Note: Accept equivalent angles, in radians or degrees.
Accept alternate answers without cis e.g. $= \frac{8a^6}{i}$

[7 marks]

4. (a)



AIAI

Note: Award *AI* for correct asymptote with correct behaviour and *AI* for shape.

[2 marks]

- (b) intersect on $y = x$
 $x + \ln x = x \Rightarrow \ln x = 0$
 intersect at (1, 1)

(MI)

(AI)

AIAI

[4 marks]

Total [6 marks]

5. (a) $\cos x = 0, \sin x = 0$ (M1)

$$x = \frac{n\pi}{2}, n \in \mathbb{Z}$$

A1

[2 marks]

- (b) **EITHER**

$$\frac{\sin 3x \cos x - \cos 3x \sin x}{\sin x \cos x}$$

M1A1

$$= \frac{\sin(3x - x)}{\frac{1}{2} \sin 2x}$$

A1A1

$$= 2$$

A1

OR

$$\frac{\sin 2x \cos x + \cos 2x \sin x}{\sin x} - \frac{\cos 2x \cos x - \sin 2x \sin x}{\cos x}$$

M1

$$= \frac{2 \sin x \cos^2 x + 2 \cos^2 x \sin x - \sin x}{\sin x} - \frac{2 \cos^3 x - \cos x - 2 \sin^2 x \cos x}{\cos x}$$

A1A1

$$= 4 \cos^2 x - 1 - 2 \cos^2 x + 1 + 2 \sin^2 x$$

A1

$$= 2 \cos^2 x + 2 \sin^2 x$$

A1

$$= 2$$

[5 marks]

Total [7 marks]

6. (a) $\int_{\frac{1}{6}}^1 \frac{k}{x} - \frac{1}{x} dx = (k-1)[\ln x]_{\frac{1}{6}}^1$ M1A1

Note: Award **M1** for $\int \frac{k}{x} - \frac{1}{x} dx$ or $\int \frac{1}{x} - \frac{k}{x} dx$ and **A1** for $(k-1) \ln x$ seen in part (a) or later in part (b).

$$= (1-k) \ln \frac{1}{6}$$

A1

[3 marks]

- (b) $\int_1^{\sqrt{6}} \frac{k}{x} - \frac{1}{x} dx = (k-1)[\ln x]_1^{\sqrt{6}}$ (A1)

Note: Award **A1** for correct change of limits.

$$= (k-1) \ln \sqrt{6}$$

A1

[2 marks]

continued ...

Question 6 continued

(c) $(1 - k)\ln \frac{1}{6} = (k - 1)\ln 6$ AI

$(k - 1)\ln \sqrt{6} = \frac{1}{2}(k - 1)\ln 6$ AI

Note: This simplification could have occurred earlier, and marks should still be awarded.

ratio is 2 (or 2:1) AI

[3 marks]

Total [8 marks]

7. $\sqrt{x^2 + y^2} + x + yi = 6 - 2i$ (AI)

equating real and imaginary parts MI

$y = -2$ AI

$\sqrt{x^2 + 4} + x = 6$ AI

$x^2 + 4 = (6 - x)^2$ MI

$-32 = -12x \Rightarrow x = \frac{8}{3}$ AI

[6 marks]

8. $\log_3 \left(\frac{9}{x+7} \right) = \log_3 \frac{1}{2x}$ MIMIAI

Note: Award **MI** for changing to single base, **MI** for incorporating the 2 into a log and **AI** for a correct equation with maximum one log expression each side.

$x + 7 = 18x$ MI

$x = \frac{7}{17}$ AI

[5 marks]

9. $4x + 2y \frac{dy}{dx} = 0 \Rightarrow \frac{dy}{dx} = -\frac{2x}{y}$ MIAI

Note: Allow follow through on incorrect $\frac{dy}{dx}$ from this point.

gradient of normal at (a, b) is $\frac{b}{2a}$

Note: No further A marks are available if a general point is not used

equation of normal at (a, b) is $y - b = \frac{b}{2a}(x - a) \left(\Rightarrow y = \frac{b}{2a}x + \frac{b}{2} \right)$ MIAI

substituting $(1, 0)$ MI

$b = 0$ or $a = -1$ AIAI

four points are $(3, 0), (-3, 0), (-1, 4), (-1, -4)$ AIAI

Note: Award **AIA0** for any two points correct.

[9 marks]

SECTION B

10.	(a)	$\cos \hat{A} = \frac{BA}{\sqrt{2}}$	<i>AI</i>
		$\sin \hat{A} = \frac{BC}{\sqrt{2}}$	<i>AI</i>
		$\cos \hat{A} - \sin \hat{A} = \frac{BA - BC}{\sqrt{2}}$	<i>RI</i>
		$= \frac{1}{\sqrt{2}}$	<i>AG</i>

[3 marks]

(b)	$\cos^2 \hat{A} - 2 \cos \hat{A} \sin \hat{A} + \sin^2 \hat{A} = \frac{1}{2}$	<i>MIAI</i>
	$1 - 2 \sin \hat{A} \cos \hat{A} = \frac{1}{2}$	<i>MIAI</i>
	$\sin 2\hat{A} = \frac{1}{2}$	<i>MI</i>
	$2\hat{A} = 30^\circ$	<i>AI</i>
	angles in the triangle are 15° and 75°	<i>AIAI</i>

Note: Accept answers in radians.

[8 marks]

(c)	$BC^2 + (BC + 1)^2 = 2$	<i>MIAI</i>
	$2BC^2 + 2BC - 1 = 0$	<i>AI</i>
	$BC = \frac{-2 + \sqrt{12}}{4} \left(= \frac{\sqrt{3} - 1}{2} \right)$	<i>MIAI</i>
	$\sin \hat{A} = \frac{BC}{\sqrt{2}} = \frac{\sqrt{3} - 1}{2\sqrt{2}}$	<i>AI</i>
	$= \frac{\sqrt{6} - \sqrt{2}}{4}$	<i>AG</i>

[6 marks]

continued ...

Question 10 continued

(d) **EITHER**

$$\begin{aligned}
 h &= AB \sin \hat{A} && \text{MI} \\
 &= (BC + 1) \sin \hat{A} && \text{AI} \\
 &= \frac{\sqrt{3} + 1}{2} \times \frac{\sqrt{6} - \sqrt{2}}{4} = \frac{\sqrt{2}}{4} && \text{MIAI}
 \end{aligned}$$

OR

$$\begin{aligned}
 \frac{1}{2} AB \cdot BC &= \frac{1}{2} AC \cdot h && \text{MI} \\
 \frac{\sqrt{3} - 1}{2} \cdot \frac{\sqrt{3} + 1}{2} &= \sqrt{2} h && \text{AI} \\
 \frac{2}{4} &= \sqrt{2} h && \text{MI} \\
 h &= \frac{1}{2\sqrt{2}} && \text{AI}
 \end{aligned}$$

[4 marks]

Total [21 marks]

11. (a) if $n = 1$

$$\begin{aligned}
 \mathbf{X}^1 &= \mathbf{U}^{-1} \mathbf{A}^1 \mathbf{U} \text{ which is given, so true for } n = 1 && \text{AI} \\
 \text{Assume true for } n = k &&& \\
 \mathbf{X}^k &= \mathbf{U}^{-1} \mathbf{A}^k \mathbf{U} && \text{MI}
 \end{aligned}$$

Note: Only award **MI** if the word “true” or equivalent appears.

$$\begin{aligned}
 \text{if } n &= k + 1 && \\
 \mathbf{X}^{k+1} &= \mathbf{X} \mathbf{X}^k && \text{MI} \\
 &= \mathbf{U}^{-1} \mathbf{A} \mathbf{U} \mathbf{U}^{-1} \mathbf{A}^k \mathbf{U} && \text{AI} \\
 &= \mathbf{U}^{-1} \mathbf{A} \mathbf{A}^k \mathbf{U} = \mathbf{U}^{-1} \mathbf{A} \mathbf{A}^k \mathbf{U} && \text{(AI)} \\
 &= \mathbf{U}^{-1} \mathbf{A}^{k+1} \mathbf{U} && \text{AI}
 \end{aligned}$$

As true for $n = 1$, and true for $n = k \Rightarrow$ true for $n = k + 1$, then by the principle of mathematical induction the statement is true for all $n (\in \mathbb{Z}^+)$ **RI**

Note: Do not award **RI** if both **M** marks have not been awarded in this part. For **RI** to be awarded evidence of implication should be seen in the statement.

[7 marks]

continued ...

Question 11 continued

(b) (i) **METHOD 1**

$$AU = UD \Rightarrow D = U^{-1}AU \quad \text{MI}$$

$$U^{-1} = \frac{1}{2} \begin{pmatrix} 1 & -1 \\ -1 & 3 \end{pmatrix} \quad \text{(AI)}$$

$$D = U^{-1}AU = \frac{1}{2} \begin{pmatrix} 1 & -1 \\ -1 & 3 \end{pmatrix} \begin{pmatrix} 2 & -3 \\ 1 & -2 \end{pmatrix} \begin{pmatrix} 3 & 1 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad \text{MIAI}$$

METHOD 2

$$\begin{pmatrix} 2 & -3 \\ 1 & -2 \end{pmatrix} \begin{pmatrix} 3 & 1 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 3 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \quad \text{MI}$$

$$\begin{pmatrix} 3 & -1 \\ 1 & -1 \end{pmatrix} = \begin{pmatrix} 3a+c & 3b+d \\ a+c & b+d \end{pmatrix} \quad \text{AI}$$

solving simultaneously MI

$$a=1, c=0, b=0, d=-1$$

$$D = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad \text{AI}$$

(ii) $D^2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \text{AI}$

(iii) $D^{2n} = U^{-1}A^{2n}U \Rightarrow A^{2n} = UD^{2n}U^{-1}$
 $\qquad \qquad \qquad = UIU^{-1}$
 $\qquad \qquad \qquad = UU^{-1} = I \quad \text{MIAI}$
AI
AG

(iv) $A^{2n} = I \Rightarrow A^n A^n = I$
 $\Rightarrow (A^n)^{-1} = A^n \quad \text{MI}$
RIAG

[10 marks]

Total [17 marks]

12. (a) **EITHER**

derivative of $\frac{x}{1-x}$ is $\frac{(1-x) - x(-1)}{(1-x)^2}$ *MIAI*

$$f'(x) = \frac{1}{2} \left(\frac{x}{1-x} \right)^{\frac{1}{2}} \frac{1}{(1-x)^2}$$
 MIAI

$$= \frac{1}{2} x^{-\frac{1}{2}} (1-x)^{-\frac{3}{2}}$$
 AG

$f'(x) > 0$ (for all $0 < x < 1$) so the function is increasing *RI*

OR

$$f(x) = \frac{x^{\frac{1}{2}}}{(1-x)^{\frac{1}{2}}}$$

$$f'(x) = \frac{(1-x)^{\frac{1}{2}} \left(\frac{1}{2} x^{-\frac{1}{2}} \right) - \frac{1}{2} x^{\frac{1}{2}} (1-x)^{-\frac{1}{2}} (-1)}{1-x}$$
 MIAI

$$= \frac{1}{2} x^{-\frac{1}{2}} (1-x)^{-\frac{1}{2}} + \frac{1}{2} x^{\frac{1}{2}} (1-x)^{-\frac{3}{2}}$$
 AI

$$= \frac{1}{2} x^{-\frac{1}{2}} (1-x)^{-\frac{3}{2}} [1-x+x]$$
 MI

$$= \frac{1}{2} x^{-\frac{1}{2}} (1-x)^{-\frac{3}{2}}$$
 AG

$f'(x) > 0$ (for all $0 < x < 1$) so the function is increasing *RI*

[5 marks]

(b) $f'(x) = \frac{1}{2} x^{-\frac{1}{2}} (1-x)^{-\frac{3}{2}}$

$$\Rightarrow f''(x) = -\frac{1}{4} x^{-\frac{3}{2}} (1-x)^{-\frac{3}{2}} + \frac{3}{4} x^{-\frac{1}{2}} (1-x)^{-\frac{5}{2}}$$
 MIAI

$$= -\frac{1}{4} x^{-\frac{3}{2}} (1-x)^{-\frac{5}{2}} [1-4x]$$

$$f''(x) = 0 \Rightarrow x = \frac{1}{4}$$
 MIAI

$f''(x)$ changes sign at $x = \frac{1}{4}$ hence there is a point of inflexion *RI*

$$x = \frac{1}{4} \Rightarrow y = \frac{1}{\sqrt{3}}$$
 AI

the coordinates are $\left(\frac{1}{4}, \frac{1}{\sqrt{3}} \right)$

[6 marks]

continued ...

Question 12 continued

(c) $x = \sin^2 \theta \Rightarrow \frac{dx}{d\theta} = 2 \sin \theta \cos \theta$ *MIAI*

$$\int \sqrt{\frac{x}{1-x}} dx = \int \sqrt{\frac{\sin^2 \theta}{1-\sin^2 \theta}} 2 \sin \theta \cos \theta d\theta$$
 MIAI

$$= \int 2 \sin^2 \theta d\theta$$
 AI

$$= \int 1 - \cos 2\theta d\theta$$
 MIAI

$$= \theta - \frac{1}{2} \sin 2\theta + c$$
 AI

$$\theta = \arcsin \sqrt{x}$$
 AI

$$\frac{1}{2} \sin 2\theta = \sin \theta \cos \theta = \sqrt{x} \sqrt{1-x} = \sqrt{x-x^2}$$
 MIAI

hence $\int \sqrt{\frac{x}{1-x}} dx = \arcsin \sqrt{x} - \sqrt{x-x^2} + c$ *AG*

[11 marks]

Total [22 marks]
